


Molecular Biology of **THE CELL**

Fifth Edition



REFERENCE EDITION

Bruce Alberts

Alexander Johnson

Julian Lewis

Martin Raff

Keith Roberts

Peter Walter

With problems by

John Wilson

Tim Hunt

Contents

| | |
|-----------------------------|---|
| <i>Special Features</i> | <i>viii</i> |
| <i>Detailed Contents</i> | <i>ix</i> |
| <i>Acknowledgments</i> | <i>xxvi</i> |
| <i>A Note to the Reader</i> | <i>xxxi</i> |
| PART I | INTRODUCTION TO THE CELL |
| 1. | Cells and Genomes 1 |
| 2. | Cell Chemistry and Biosynthesis 45 |
| 3. | Proteins 125 |
| PART II | BASIC GENETIC MECHANISMS |
| 4. | DNA, Chromosomes, and Genomes 195 |
| 5. | DNA Replication, Repair, and Recombination 263 |
| 6. | How Cells Read the Genome: From DNA to Protein 329 |
| 7. | Control of Gene Expression 411 |
| PART III | METHODS |
| 8. | Manipulating Proteins, DNA, and RNA 501 |
| 9. | Visualizing Cells 579 |
| PART IV | INTERNAL ORGANIZATION OF THE CELL |
| 10. | Membrane Structure 617 |
| 11. | Membrane Transport of Small Molecules and the Electrical Properties of Membranes 651 |
| 12. | Intracellular Compartments and Protein Sorting 695 |
| 13. | Intracellular Vesicular Traffic 749 |
| 14. | Energy Conversion: Mitochondria and Chloroplasts 813 |
| 15. | Mechanisms of Cell Communication 879 |
| 16. | The Cytoskeleton 965 |
| 17. | The Cell Cycle 1053 |
| 18. | Apoptosis 1115 |
| PART V | CELLS IN THEIR SOCIAL CONTEXT |
| 19. | Cell Junctions, Cell Adhesion, and the Extracellular Matrix 1131 |
| 20. | Cancer 1205 |
| 21. | Sexual Reproduction: Meiosis, Germ Cells, and Fertilization 1269 |
| 22. | Development of Multicellular Organisms 1305 |
| 23. | Specialized Tissues, Stem Cells, and Tissue Renewal 1417 |
| 24. | Pathogens, Infection, and Innate Immunity 1485 |
| 25. | The Adaptive Immune System 1539 |
| <i>Glossary</i> | <i>G-1</i> |
| <i>Index</i> | <i>I-1</i> |
| <i>Tables</i> | <i>The Genetic Code, Amino Acids T-1</i> |

Detailed Contents

Chapter 1 Cells and Genomes

THE UNIVERSAL FEATURES OF CELLS ON EARTH

| | |
|---|----|
| All Cells Store Their Hereditary Information in the Same Linear Chemical Code (DNA) | 1 |
| All Cells Replicate Their Hereditary Information by Templated Polymerization | 2 |
| All Cells Transcribe Portions of Their Hereditary Information into the Same Intermediary Form (RNA) | 3 |
| All Cells Use Proteins as Catalysts | 4 |
| All Cells Translate RNA into Protein in the Same Way | 5 |
| The Fragment of Genetic Information Corresponding to One Protein Is One Gene | 6 |
| Life Requires Free Energy | 7 |
| All Cells Function as Biochemical Factories Dealing with the Same Basic Molecular Building Blocks | 8 |
| All Cells Are Enclosed in a Plasma Membrane Across Which Nutrients and Waste Materials Must Pass | 9 |
| A Living Cell Can Exist with Fewer Than 500 Genes | 10 |
| Summary | 11 |

THE DIVERSITY OF GENOMES AND THE TREE OF LIFE

| | |
|--|----|
| Cells Can Be Powered by a Variety of Free Energy Sources | 11 |
| Some Cells Fix Nitrogen and Carbon Dioxide for Others | 12 |
| The Greatest Biochemical Diversity Exists Among Prokaryotic Cells | 13 |
| The Tree of Life Has Three Primary Branches: Bacteria, Archaea, and Eucaryotes | 14 |
| Some Genes Evolve Rapidly; Others Are Highly Conserved | 15 |
| Most Bacteria and Archaea Have 1000–6000 Genes | 16 |
| New Genes Are Generated from Preexisting Genes | 17 |
| Gene Duplications Give Rise to Families of Related Genes Within a Single Cell | 18 |
| Genes Can Be Transferred Between Organisms, Both in the Laboratory and in Nature | 19 |
| Sex Results in Horizontal Exchanges of Genetic Information Within a Species | 21 |
| The Function of a Gene Can Often Be Deduced from Its Sequence | 22 |
| More Than 200 Gene Families Are Common to All Three Primary Branches of the Tree of Life | 22 |
| Mutations Reveal the Functions of Genes | 23 |
| Molecular Biologists Have Focused a Spotlight on <i>E. coli</i> | 23 |
| Summary | 24 |

GENETIC INFORMATION IN EUCARYOTES

| | |
|---|----|
| Eucaryotic Cells May Have Originated as Predators | 26 |
| Modern Eucaryotic Cells Evolved from a Symbiosis | 26 |
| Eucaryotes Have Hybrid Genomes | 27 |
| Eucaryotic Genomes Are Big | 30 |
| Eucaryotic Genomes Are Rich in Regulatory DNA | 30 |
| The Genome Defines the Program of Multicellular Development | 31 |
| Many Eucaryotes Live as Solitary Cells: the Protists | 31 |
| A Yeast Serves as a Minimal Model Eucaryote | 32 |
| The Expression Levels of All The Genes of An Organism Can Be Monitored Simultaneously | 33 |
| To Make Sense of Cells, We Need Mathematics, Computers, and Quantitative Information | 34 |
| <i>Arabidopsis</i> Has Been Chosen Out of 300,000 Species As a Model Plant | 35 |
| | 36 |

| | |
|--|----|
| The World of Animal Cells Is Represented By a Worm, a Fly, a Mouse, and a Human | 36 |
| Studies in <i>Drosophila</i> Provide a Key to Vertebrate Development | 37 |
| The Vertebrate Genome Is a Product of Repeated Duplication | 38 |
| Genetic Redundancy Is a Problem for Geneticists, But It Creates Opportunities for Evolving Organisms | 39 |
| The Mouse Serves as a Model for Mammals | 39 |
| Humans Report on Their Own Peculiarities | 40 |
| We Are All Different in Detail | 41 |
| Summary | 42 |
| Problems | 42 |
| References | 44 |

Chapter 2 Cell Chemistry and Biosynthesis 45

THE CHEMICAL COMPONENTS OF A CELL 45

| | |
|--|----|
| Cells Are Made From a Few Types of Atoms | 45 |
| The Outermost Electrons Determine How Atoms Interact | 46 |
| Covalent Bonds Form by the Sharing of Electrons | 48 |
| There Are Different Types of Covalent Bonds | 50 |
| An Atom Often Behaves as if It Has a Fixed Radius | 51 |
| Water Is the Most Abundant Substance in Cells | 51 |
| Some Polar Molecules Are Acids and Bases | 52 |
| Four Types of Noncovalent Attractions Help Bring Molecules Together in Cells | 53 |
| A Cell Is Formed from Carbon Compounds | 54 |
| Cells Contain Four Major Families of Small Organic Molecules | 55 |
| Sugars Provide an Energy Source for Cells and Are the Subunits of Polysaccharides | 55 |
| Fatty Acids Are Components of Cell Membranes, as Well as a Source of Energy | 58 |
| Amino Acids Are the Subunits of Proteins | 59 |
| Nucleotides Are the Subunits of DNA and RNA | 61 |
| The Chemistry of Cells Is Dominated by Macromolecules with Remarkable Properties | 62 |
| Noncovalent Bonds Specify Both the Precise Shape of a Macromolecule and its Binding to Other Molecules | 63 |
| Summary | 65 |

CATALYSIS AND THE USE OF ENERGY BY CELLS 65

| | |
|---|----|
| Cell Metabolism Is Organized by Enzymes | 66 |
| Biological Order Is Made Possible by the Release of Heat Energy from Cells | 66 |
| Photosynthetic Organisms Use Sunlight to Synthesize Organic Molecules | 68 |
| Cells Obtain Energy by the Oxidation of Organic Molecules | 70 |
| Oxidation and Reduction Involve Electron Transfers | 71 |
| Enzymes Lower the Barriers That Block Chemical Reactions | 72 |
| How Enzymes Find Their Substrates: The Enormous Rapidity of Molecular Motions | 74 |
| The Free-Energy Change for a Reaction Determines Whether It Can Occur | 75 |
| The Concentration of Reactants Influences the Free-Energy Change and a Reaction's Direction | 76 |
| For Sequential Reactions, ΔG° Values Are Additive | 77 |
| Activated Carrier Molecules Are Essential for Biosynthesis | 78 |
| The Formation of an Activated Carrier Is Coupled to an Energetically Favorable Reaction | 79 |

| | | | |
|--|------------|---|------------|
| ATP Is the Most Widely Used Activated Carrier Molecule | 80 | Molecular Tunnels Channel Substrates in Enzymes with Multiple Catalytic Sites | 167 |
| Energy Stored in ATP Is Often Harnessed to Join Two Molecules Together | 81 | Multienzyme Complexes Help to Increase the Rate of Cell Metabolism | 168 |
| NADH and NADPH Are Important Electron Carriers | 82 | The Cell Regulates the Catalytic Activities of its Enzymes | 169 |
| There Are Many Other Activated Carrier Molecules in Cells | 83 | Allosteric Enzymes Have Two or More Binding Sites That Interact | 171 |
| The Synthesis of Biological Polymers Is Driven by ATP Hydrolysis | 84 | Two Ligands Whose Binding Sites Are Coupled Must Reciprocally Affect Each Other's Binding | 171 |
| <i>Summary</i> | 87 | Symmetric Protein Assemblies Produce Cooperative Allosteric Transitions | 172 |
| HOW CELLS OBTAIN ENERGY FROM FOOD | 88 | The Allosteric Transition in Aspartate Transcarbamoylase Is Understood in Atomic Detail | 173 |
| Glycolysis Is a Central ATP-Producing Pathway | 88 | Many Changes in Proteins Are Driven by Protein Phosphorylation | 175 |
| Fermentations Produce ATP in the Absence of Oxygen | 89 | A Eucaryotic Cell Contains a Large Collection of Protein Kinases and Protein Phosphatases | 176 |
| Glycolysis Illustrates How Enzymes Couple Oxidation to Energy Storage | 91 | The Regulation of Cdk and Src Protein Kinases Shows How a Protein Can Function as a Microchip | 177 |
| Organisms Store Food Molecules in Special Reservoirs | 91 | Proteins That Bind and Hydrolyze GTP Are Ubiquitous Cellular Regulators | 178 |
| Most Animal Cells Derive Their Energy from Fatty Acids Between Meals | 95 | Regulatory Proteins Control the Activity of GTP-Binding Proteins by Determining Whether GTP or GDP Is Bound | 179 |
| Sugars and Fats Are Both Degraded to Acetyl CoA in Mitochondria | 96 | Large Protein Movements Can Be Generated From Small Ones | 179 |
| The Citric Acid Cycle Generates NADH by Oxidizing Acetyl Groups to CO ₂ | 97 | Motor Proteins Produce Large Movements in Cells | 181 |
| Electron Transport Drives the Synthesis of the Majority of the ATP in Most Cells | 100 | Membrane-Bound Transporters Harness Energy to Pump Molecules Through Membranes | 182 |
| Amino Acids and Nucleotides Are Part of the Nitrogen Cycle | 100 | Proteins Often Form Large Complexes That Function as Protein Machines | 184 |
| Metabolism Is Organized and Regulated | 101 | Protein Machines with Interchangeable Parts Make Efficient Use of Genetic Information | 184 |
| <i>Summary</i> | 103 | The Activation of Protein Machines Often Involves Positioning Them at Specific Sites | 185 |
| <i>Problems</i> | 103 | Many Proteins Are Controlled by Multisite Covalent Modification | 186 |
| <i>References</i> | 124 | A Complex Network of Protein Interactions Underlies Cell Function | 187 |
| Chapter 3 Proteins | 125 | <i>Summary</i> | 190 |
| THE SHAPE AND STRUCTURE OF PROTEINS | 125 | <i>Problems</i> | 191 |
| The Shape of a Protein Is Specified by Its Amino Acid Sequence | 125 | <i>References</i> | 193 |
| Proteins Fold into a Conformation of Lowest Energy | 130 | Chapter 4 DNA, Chromosomes, and Genomes | 195 |
| The α Helix and the β Sheet Are Common Folding Patterns | 131 | THE STRUCTURE AND FUNCTION OF DNA | 197 |
| Protein Domains Are Modular Units from which Larger Proteins Are Built | 135 | A DNA Molecule Consists of Two Complementary Chains of Nucleotides | 197 |
| Few of the Many Possible Polypeptide Chains Will Be Useful to Cells | 136 | The Structure of DNA Provides a Mechanism for Heredity | 199 |
| Proteins Can Be Classified into Many Families | 137 | In Eucaryotes, DNA Is Enclosed in a Cell Nucleus | 200 |
| Sequence Searches Can Identify Close Relatives | 139 | <i>Summary</i> | 201 |
| Some Protein Domains Form Parts of Many Different Proteins | 140 | CHROMOSOMAL DNA AND ITS PACKAGING IN THE CHROMATIN FIBER | 202 |
| Certain Pairs of Domains Are Found Together in Many Proteins | 141 | Eucaryotic DNA Is Packaged into a Set of Chromosomes | 202 |
| The Human Genome Encodes a Complex Set of Proteins, Revealing Much That Remains Unknown | 142 | Chromosomes Contain Long Strings of Genes | 204 |
| Larger Protein Molecules Often Contain More Than One Polypeptide Chain | 142 | The Nucleotide Sequence of the Human Genome Shows How Our Genes Are Arranged | 205 |
| Some Proteins Form Long Helical Filaments | 143 | Genome Comparisons Reveal Evolutionarily Conserved DNA Sequences | 207 |
| Many Protein Molecules Have Elongated, Fibrous Shapes | 145 | Chromosomes Exist in Different States Throughout the Life of a Cell | 208 |
| Many Proteins Contain a Surprisingly Large Amount of Unstructured Polypeptide Chain | 146 | Each DNA Molecule That Forms a Linear Chromosome Must Contain a Centromere, Two Telomeres, and Replication Origins | 209 |
| Covalent Cross-Linkages Often Stabilize Extracellular Proteins | 147 | DNA Molecules Are Highly Condensed in Chromosomes | 210 |
| Protein Molecules Often Serve as Subunits for the Assembly of Large Structures | 148 | Nucleosomes Are a Basic Unit of Eucaryotic Chromosome Structure | 211 |
| Many Structures in Cells Are Capable of Self-Assembly | 149 | The Structure of the Nucleosome Core Particle Reveals How DNA Is Packaged | 212 |
| Assembly Factors Often Aid the Formation of Complex Biological Structures | 151 | Nucleosomes Have a Dynamic Structure, and Are Frequently Subjected to Changes Catalyzed by ATP-Dependent Chromatin-Remodeling Complexes | 215 |
| <i>Summary</i> | 152 | Nucleosomes Are Usually Packed Together into a Compact Chromatin Fiber | 216 |
| PROTEIN FUNCTION | 152 | <i>Summary</i> | 218 |
| All Proteins Bind to Other Molecules | 153 | THE REGULATION OF CHROMATIN STRUCTURE | 219 |
| The Surface Conformation of a Protein Determines Its Chemistry | 154 | Some Early Mysteries Concerning Chromatin Structure | 220 |
| Sequence Comparisons Between Protein Family Members Highlight Crucial Ligand-Binding Sites | 155 | | |
| Proteins Bind to Other Proteins Through Several Types of Interfaces | 156 | | |
| Antibody Binding Sites Are Especially Versatile | 156 | | |
| The Equilibrium Constant Measures Binding Strength | 157 | | |
| Enzymes Are Powerful and Highly Specific Catalysts | 158 | | |
| Substrate Binding Is the First Step in Enzyme Catalysis | 159 | | |
| Enzymes Speed Reactions by Selectively Stabilizing Transition States | 160 | | |
| Enzymes Can Use Simultaneous Acid and Base Catalysis | 160 | | |
| Lysozyme Illustrates How an Enzyme Works | 161 | | |
| Tightly Bound Small Molecules Add Extra Functions to Proteins | 166 | | |

| | | | |
|--|-----|---|-----|
| Heterochromatin Is Highly Organized and Unusually Resistant to Gene Expression | 220 | DNA REPLICATION MECHANISMS | 266 |
| The Core Histones Are Covalently Modified at Many Different Sites | 222 | Base-Pairing Underlies DNA Replication and DNA Repair | 266 |
| Chromatin Acquires Additional Variety through the Site-Specific Insertion of a Small Set of Histone Variants | 224 | The DNA Replication Fork Is Asymmetrical | 266 |
| The Covalent Modifications and the Histone Variants Act in Concert to Produce a "Histone Code" That Helps to Determine Biological Function | 224 | The High Fidelity of DNA Replication Requires Several Proofreading Mechanisms | 268 |
| A Complex of Code-Reader and Code-Writer Proteins Can Spread Specific Chromatin Modifications for Long Distances Along a Chromosome | 226 | Only DNA Replication in the 5'-to-3' Direction Allows Efficient Error Correction | 271 |
| Barrier DNA Sequences Block the Spread of Reader-Writer Complexes and Thereby Separate Neighboring Chromatin Domains | 227 | A Special Nucleotide-Polymerizing Enzyme Synthesizes Short RNA Primer Molecules on the Lagging Strand | 272 |
| The Chromatin in Centromeres Reveals How Histone Variants Can Create Special Structures | 228 | Special Proteins Help to Open Up the DNA Double Helix in Front of the Replication Fork | 273 |
| Chromatin Structures Can Be Directly Inherited | 230 | A Sliding Ring Holds a Moving DNA Polymerase onto the DNA | 273 |
| Chromatin Structures Add Unique Features to Eucaryotic Chromosome Function | 231 | The Proteins at a Replication Fork Cooperate to Form a Replication Machine | 275 |
| <i>Summary</i> | 233 | A Strand-Directed Mismatch Repair System Removes Replication Errors That Escape from the Replication Machine | 276 |
| THE GLOBAL STRUCTURE OF CHROMOSOMES | 233 | DNA Topoisomerases Prevent DNA Tangling During Replication | 278 |
| Chromosomes Are Folded into Large Loops of Chromatin | 234 | DNA Replication Is Fundamentally Similar in Eucaryotes and Bacteria | 280 |
| Polytene Chromosomes Are Uniquely Useful for Visualizing Chromatin Structures | 236 | <i>Summary</i> | 281 |
| There Are Multiple Forms of Heterochromatin | 238 | THE INITIATION AND COMPLETION OF DNA REPLICATION IN CHROMOSOMES | 281 |
| Chromatin Loops Decondense When the Genes Within Them Are Expressed | 239 | DNA Synthesis Begins at Replication Origins | 281 |
| Chromatin Can Move to Specific Sites Within the Nucleus to Alter Their Gene Expression | 239 | Bacterial Chromosomes Typically Have a Single Origin of DNA Replication | 282 |
| Networks of Macromolecules Form a Set of Distinct Biochemical Environments inside the Nucleus | 241 | Eucaryotic Chromosomes Contain Multiple Origins of Replication | 282 |
| Mitotic Chromosomes Are Formed from Chromatin in Its Most Condensed State | 243 | In Eucaryotes DNA Replication Takes Place During Only One Part of the Cell Cycle | 284 |
| <i>Summary</i> | 245 | Different Regions on the Same Chromosome Replicate at Distinct Times in S Phase | 285 |
| HOW GENOMES EVOLVE | 245 | Highly Condensed Chromatin Replicates Late, While Genes in Less Condensed Chromatin Tend to Replicate Early | 285 |
| Genome Alterations Are Caused by Failures of the Normal Mechanisms for Copying and Maintaining DNA | 246 | Well-Defined DNA Sequences Serve as Replication Origins in a Simple Eucaryote, the Budding Yeast | 286 |
| The Genome Sequences of Two Species Differ in Proportion to the Length of Time That They Have Separately Evolved | 247 | A Large Multisubunit Complex Binds to Eucaryotic Origins of Replication | 287 |
| Phylogenetic Trees Constructed from a Comparison of DNA Sequences Trace the Relationships of All Organisms | 248 | The Mammalian DNA Sequences That Specify the Initiation of Replication Have Been Difficult to Identify | 288 |
| A Comparison of Human and Mouse Chromosomes Shows How the Structures of Genomes Diverge | 249 | New Nucleosomes Are Assembled Behind the Replication Fork | 289 |
| The Size of a Vertebrate Genome Reflects the Relative Rates of DNA Addition and DNA Loss in a Lineage | 251 | The Mechanisms of Eucaryotic Chromosome Duplication Ensure That Patterns of Histone Modification Can Be Inherited | 290 |
| We Can Reconstruct the Sequence of Some Ancient Genomes | 251 | Telomerase Replicates the Ends of Chromosomes | 292 |
| Multispecies Sequence Comparisons Identify Important DNA Sequences of Unknown Function | 252 | Telomere Length Is Regulated by Cells and Organisms | 293 |
| Accelerated Changes in Previously Conserved Sequences Can Help Decipher Critical Steps in Human Evolution | 253 | <i>Summary</i> | 294 |
| Gene Duplication Provides an Important Source of Genetic Novelty During Evolution | 253 | DNA REPAIR | 295 |
| Duplicated Genes Diverge | 254 | Without DNA Repair, Spontaneous DNA Damage Would Rapidly Change DNA Sequences | 296 |
| The Evolution of the Globin Gene Family Shows How DNA Duplications Contribute to the Evolution of Organisms | 256 | The DNA Double Helix Is Readily Repaired | 296 |
| Genes Encoding New Proteins Can Be Created by the Recombination of Exons | 257 | DNA Damage Can Be Removed by More Than One Pathway | 297 |
| Neutral Mutations Often Spread to Become Fixed in a Population, with a Probability that Depends on Population Size | 257 | Coupling DNA Repair to Transcription Ensures That the Cell's Most Important DNA Is Efficiently Repaired | 299 |
| A Great Deal Can Be Learned from Analyses of the Variation Among Humans | 258 | The Chemistry of the DNA Bases Facilitates Damage Detection | 300 |
| <i>Summary</i> | 260 | Special DNA Polymerases Are Used in Emergencies to Repair DNA | 302 |
| <i>Problems</i> | 260 | Double-Strand Breaks Are Efficiently Repaired | 302 |
| <i>References</i> | 262 | DNA Damage Delays Progression of the Cell Cycle | 303 |
| | | <i>Summary</i> | 304 |
| Chapter 5 DNA Replication, Repair, and Recombination | 263 | HOMOLOGOUS RECOMBINATION | 304 |
| THE MAINTENANCE OF DNA SEQUENCES | 263 | Homologous Recombination Has Many Uses in the Cell | 304 |
| Mutation Rates Are Extremely Low | 263 | Homologous Recombination Has Common Features in All Cells | 305 |
| Low Mutation Rates Are Necessary for Life as We Know It | 265 | DNA Base-Pairing Guides Homologous Recombination | 305 |
| <i>Summary</i> | 265 | The RecA Protein and its Homologs Enable a DNA Single Strand to Pair with a Homologous Region of DNA Double Helix | 307 |
| | | Branch Migration Can Either Enlarge Heteroduplex Regions or Release Newly Synthesized DNA as a Single Strand | 308 |
| | | Homologous Recombination Can Flawlessly Repair Double-Stranded Breaks in DNA | 308 |
| | | Cells Carefully Regulate the Use of Homologous Recombination in DNA Repair | 310 |
| | | Holliday Junctions Are Often Formed During Homologous Recombination Events | 311 |

| | | | |
|---|-----|---|-----|
| Meiotic Recombination Begins with a Programmed Double-Strand Break | 312 | FROM RNA TO PROTEIN | 366 |
| Homologous Recombination Often Results in Gene Conversion | 314 | An mRNA Sequence Is Decoded in Sets of Three Nucleotide | 367 |
| Mismatch Proofreading Prevents Promiscuous Recombination Between Two Poorly Matched DNA Sequences | 315 | tRNA Molecules Match Amino Acids to Codons in mRNA | 368 |
| Summary | 316 | tRNAs Are Covalently Modified Before They Exit from the Nucleus | 369 |
| TRANSPOSITION AND CONSERVATIVE SITE-SPECIFIC RECOMBINATION | 316 | Specific Enzymes Couple Each Amino Acid to Its Appropriate tRNA Molecule | 370 |
| Through Transposition, Mobile Genetic Elements Can Insert Into Any DNA Sequence | 317 | Editing by RNA Synthetases Ensures Accuracy | 371 |
| DNA-Only Transposons Move by Both Cut-and-Paste and Replicative Mechanisms | 317 | Amino Acids Are Added to the C-terminal End of a Growing Polypeptide Chain | 373 |
| Some Viruses Use a Transposition Mechanism to Move Themselves into Host Cell Chromosomes | 319 | The RNA Message Is Decoded in Ribosomes | 373 |
| Retroviral-like Retrotransposons Resemble Retroviruses, but Lack a Protein Coat | 320 | Elongation Factors Drive Translation Forward and Improve Its Accuracy | 377 |
| A Large Fraction of the Human Genome Is Composed of Nonretroviral Retrotransposons | 321 | The Ribosome Is a Ribozyme | 378 |
| Different Transposable Elements Predominate in Different Organisms | 322 | Nucleotide Sequences in mRNA Signal Where to Start Protein Synthesis | 379 |
| Genome Sequences Reveal the Approximate Times that Transposable Elements Have Moved | 323 | Stop Codons Mark the End of Translation | 381 |
| Conservative Site-Specific Recombination Can Reversibly Rearrange DNA | 323 | Proteins Are Made on Polyribosomes | 381 |
| Conservative Site-Specific Recombination Was Discovered in Bacteriophage λ | 324 | There Are Minor Variations in the Standard Genetic Code | 382 |
| Conservative Site-Specific Recombination Can Be Used to Turn Genes On or Off | 324 | Inhibitors of Prokaryotic Protein Synthesis Are Useful as Antibiotics | 383 |
| Summary | 326 | Accuracy in Translation Requires the Expenditure of Free Energy | 385 |
| Problems | 327 | Quality Control Mechanisms Act to Prevent Translation of Damaged mRNAs | 385 |
| References | 328 | Some Proteins Begin to Fold While Still Being Synthesized | 387 |
| Chapter 6 How Cells Read the Genome: From DNA to Protein | 329 | Molecular Chaperones Help Guide the Folding of Most Proteins | 388 |
| FROM DNA TO RNA | 331 | Exposed Hydrophobic Regions Provide Critical Signals for Protein Quality Control | 390 |
| Portions of DNA Sequence Are Transcribed into RNA | 332 | The Proteasome Is a Compartmentalized Protease with Sequestered Active Sites | 391 |
| Transcription Produces RNA Complementary to One Strand of DNA | 333 | An Elaborate Ubiquitin-Conjugating System Marks Proteins for Destruction | 393 |
| Cells Produce Several Types of RNA | 335 | Many Proteins Are Controlled by Regulated Destruction | 395 |
| Signals Encoded in DNA Tell RNA Polymerase Where to Start and Stop | 336 | Abnormally Folded Proteins Can Aggregate to Cause Destructive Human Diseases | 396 |
| Transcription Start and Stop Signals Are Heterogeneous in Nucleotide Sequence | 338 | There Are Many Steps From DNA to Protein | 399 |
| Transcription Initiation in Eucaryotes Requires Many Proteins | 339 | Summary | 399 |
| RNA Polymerase II Requires General Transcription Factors | 340 | THE RNA WORLD AND THE ORIGINS OF LIFE | 400 |
| Polymerase II Also Requires Activator, Mediator, and Chromatin-Modifying Proteins | 342 | Life Requires Stored Information | 401 |
| Transcription Elongation Produces Superhelical Tension in DNA | 343 | Polynucleotides Can Both Store Information and Catalyze Chemical Reactions | 401 |
| Transcription Elongation in Eucaryotes Is Tightly Coupled to RNA Processing | 345 | A Pre-RNA World May Predate the RNA World | 402 |
| RNA Capping Is the First Modification of Eucaryotic Pre-mRNAs | 346 | Single-Stranded RNA Molecules Can Fold into Highly Elaborate Structures | 403 |
| RNA Splicing Removes Intron Sequences from Newly Transcribed Pre-mRNAs | 347 | Self-Replicating Molecules Undergo Natural Selection | 404 |
| Nucleotide Sequences Signal Where Splicing Occurs | 349 | How Did Protein Synthesis Evolve? | 407 |
| RNA Splicing Is Performed by the Spliceosome | 349 | All Present-Day Cells Use DNA as Their Hereditary Material | 408 |
| The Spliceosome Uses ATP Hydrolysis to Produce a Complex Series of RNA-RNA Rearrangements | 351 | Summary | 408 |
| Other Properties of Pre-mRNA and Its Synthesis Help to Explain the Choice of Proper Splice Sites | 352 | Problems | 409 |
| A Second Set of snRNPs Splice a Small Fraction of Intron Sequences in Animals and Plants | 353 | References | 410 |
| RNA Splicing Shows Remarkable Plasticity | 355 | Chapter 7 Control of Gene Expression | 411 |
| Spliceosome-Catalyzed RNA Splicing Probably Evolved from Self-Splicing Mechanisms | 355 | AN OVERVIEW OF GENE CONTROL | 411 |
| RNA-Processing Enzymes Generate the 3' End of Eucaryotic mRNAs | 357 | The Different Cell Types of a Multicellular Organism Contain the Same DNA | 411 |
| Mature Eucaryotic mRNAs Are Selectively Exported from the Nucleus | 358 | Different Cell Types Synthesize Different Sets of Proteins | 412 |
| Many Noncoding RNAs Are Also Synthesized and Processed in the Nucleus | 360 | External Signals Can Cause a Cell to Change the Expression of Its Genes | 413 |
| The Nucleolus Is a Ribosome-Producing Factory | 362 | Gene Expression Can Be Regulated at Many of the Steps in the Pathway from DNA to RNA to Protein | 415 |
| The Nucleus Contains a Variety of Subnuclear Structures | 363 | Summary | 415 |
| Summary | 366 | DNA-BINDING MOTIFS IN GENETIC REGULATORY PROTEINS | 416 |
| | | Gene Regulatory Proteins Were Discovered Using Bacterial Genetics | 416 |
| | | The Outside of the DNA Helix Can Be Read by Proteins | 416 |
| | | Short DNA Sequences Are Fundamental Components of Genetic Switches | 418 |
| | | Gene Regulatory Proteins Contain Structural Motifs That Can Read DNA Sequences | 418 |
| | | The Helix-Turn-Helix Motif Is One of the Simplest and Most Common DNA-Binding Motifs | 419 |

| | | | |
|--|-----|--|-----|
| Homeodomain Proteins Constitute a Special Class of Helix–Turn–Helix Proteins | 420 | Expression of a Critical Gene Regulatory Protein Can Trigger the Expression of a Whole Battery of Downstream Genes | 463 |
| There Are Several Types of DNA-Binding Zinc Finger Motifs | 421 | Combinatorial Gene Control Creates Many Different Cell Types in Eucaryotes | 464 |
| β sheets Can Also Recognize DNA | 422 | A Single Gene Regulatory Protein Can Trigger the Formation of an Entire Organ | 465 |
| Some Proteins Use Loops That Enter the Major and Minor Groove to Recognize DNA | 423 | The Pattern of DNA Methylation Can Be Inherited When Vertebrate Cells Divide | 467 |
| The Leucine Zipper Motif Mediates Both DNA Binding and Protein Dimerization | 423 | Genomic Imprinting Is Based on DNA Methylation | 468 |
| Heterodimerization Expands the Repertoire of DNA Sequences That Gene Regulatory Proteins Can Recognize | 424 | CG-Rich Islands Are Associated with Many Genes in Mammals | 470 |
| The Helix–Loop–Helix Motif Also Mediates Dimerization and DNA Binding | 425 | Epigenetic Mechanisms Ensure That Stable Patterns of Gene Expression Can Be Transmitted to Daughter Cells | 471 |
| It Is Not Yet Possible to Predict the DNA Sequences Recognized by All Gene Regulatory Proteins | 426 | Chromosome-Wide Alterations in Chromatin Structure Can Be Inherited | 473 |
| A Gel-Mobility Shift Assay Readily Detects Sequence-Specific DNA-Binding Proteins | 427 | The Control of Gene Expression is Intrinsically Noisy | 476 |
| DNA Affinity Chromatography Facilitates the Purification of Sequence-Specific DNA-Binding Proteins | 428 | Summary | 477 |
| The DNA Sequence Recognized by a Gene Regulatory Protein Can Be Determined Experimentally | 429 | POST-TRANSCRIPTIONAL CONTROLS | 477 |
| Phylogenetic Footprinting Identifies DNA Regulatory Sequences Through Comparative Genomics | 431 | Transcription Attenuation Causes the Premature Termination of Some RNA Molecules | 477 |
| Chromatin Immunoprecipitation Identifies Many of the Sites That Gene Regulatory Proteins Occupy in Living Cells | 431 | Riboswitches Might Represent Ancient Forms of Gene Control | 478 |
| Summary | 432 | Alternative RNA Splicing Can Produce Different Forms of a Protein from the Same Gene | 479 |
| HOW GENETIC SWITCHES WORK | 432 | The Definition of a Gene Has Had to Be Modified Since the Discovery of Alternative RNA Splicing | 480 |
| The Tryptophan Repressor Is a Simple Switch That Turns Genes On and Off in Bacteria | 433 | Sex Determination in <i>Drosophila</i> Depends on a Regulated Series of RNA Splicing Events | 481 |
| Transcriptional Activators Turn Genes On | 435 | A Change in the Site of RNA Transcript Cleavage and Poly-A Addition Can Change the C-terminus of a Protein | 482 |
| A Transcriptional Activator and a Transcriptional Repressor Control the <i>Lac</i> Operon | 435 | RNA Editing Can Change the Meaning of the RNA Message | 483 |
| DNA Looping Occurs During Bacterial Gene Regulation | 437 | RNA Transport from the Nucleus Can Be Regulated | 485 |
| Bacteria Use Interchangeable RNA Polymerase Subunits to Help Regulate Gene Transcription | 438 | Some mRNAs Are Localized to Specific Regions of the Cytoplasm | 486 |
| Complex Switches Have Evolved to Control Gene Transcription in Eucaryotes | 439 | The 5' and 3' Untranslated Regions of mRNAs Control Their Translation | 488 |
| A Eucaryotic Gene Control Region Consists of a Promoter Plus Regulatory DNA Sequences | 440 | The Phosphorylation of an Initiation Factor Regulates Protein Synthesis Globally | 488 |
| Eucaryotic Gene Activator Proteins Promote the Assembly of RNA Polymerase and the General Transcription Factors at the Startpoint of Transcription | 441 | Initiation at AUG Codons Upstream of the Translation Start Can Regulate Eucaryotic Translation Initiation | 489 |
| Eucaryotic Gene Activator Proteins Also Modify Local Chromatin Structure | 442 | Internal Ribosome Entry Sites Provide Opportunities for Translation Control | 491 |
| Gene Activator Proteins Work Synergistically | 444 | Changes in mRNA Stability Can Regulate Gene Expression | 492 |
| Eucaryotic Gene Repressor Proteins Can Inhibit Transcription in Various Ways | 445 | Cytoplasmic Poly-A Addition Can Regulate Translation | 493 |
| Eucaryotic Gene Regulatory Proteins Often Bind DNA Cooperatively | 445 | Small Noncoding RNA Transcripts Regulate Many Animal and Plant Genes | 493 |
| Complex Genetic Switches That Regulate <i>Drosophila</i> Development Are Built Up from Smaller Modules | 447 | RNA Interference Is a Cell Defense Mechanism | 495 |
| The <i>Drosophila Eve</i> Gene Is Regulated by Combinatorial Controls | 448 | RNA Interference Can Direct Heterochromatin Formation | 496 |
| Complex Mammalian Gene Control Regions Are Also Constructed from Simple Regulatory Modules | 450 | RNA Interference Has Become a Powerful Experimental Tool | 497 |
| Insulators Are DNA Sequences That Prevent Eucaryotic Gene Regulatory Proteins from Influencing Distant Genes | 452 | Summary | 497 |
| Gene Switches Rapidly Evolve | 453 | Problems | 497 |
| Summary | 453 | References | 499 |
| THE MOLECULAR GENETIC MECHANISMS THAT CREATE SPECIALIZED CELL TYPES | 454 | Chapter 8 Manipulating Proteins, DNA, and RNA | 501 |
| DNA Rearrangements Mediate Phase Variation in Bacteria | 454 | ISOLATING CELLS AND GROWING THEM IN CULTURE | 501 |
| A Set of Gene Regulatory Proteins Determines Cell Type in a Budding Yeast | 455 | Cells Can Be Isolated from Intact Tissues | 502 |
| Two Proteins That Repress Each Other's Synthesis Determine the Heritable State of Bacteriophage Lambda | 457 | Cells Can Be Grown in Culture | 502 |
| Simple Gene Regulatory Circuits Can Be Used to Make Memory Devices | 458 | Eucaryotic Cell Lines Are a Widely Used Source of Homogeneous Cells | 505 |
| Transcriptional Circuits Allow the Cell to Carry Out Logic Operations | 459 | Embryonic Stem Cells Could Revolutionize Medicine | 505 |
| Synthetic Biology Creates New Devices from Existing Biological Parts | 460 | Somatic Cell Nuclear Transplantation May Provide a Way to Generate Personalized Stem Cells | 507 |
| Circadian Clocks Are Based on Feedback Loops in Gene Regulation | 460 | Hybridoma Cell Lines Are Factories That Produce Monoclonal Antibodies | 508 |
| A Single Gene Regulatory Protein Can Coordinate the Expression of a Set of Genes | 462 | Summary | 510 |
| | | PURIFYING PROTEINS | 510 |
| | | Cells Can Be Separated into Their Component Fractions | 510 |
| | | Cell Extracts Provide Accessible Systems to Study Cell Functions | 511 |
| | | Proteins Can Be Separated by Chromatography | 512 |
| | | Affinity Chromatography Exploits Specific Binding Sites on Proteins | 513 |
| | | Genetically-Engineered Tags Provide an Easy Way to Purify Proteins | 514 |

| | | | |
|---|-----|---|------------|
| Purified Cell-Free Systems Are Required for the Precise Dissection of Molecular Functions | 516 | Large Collections of Tagged Knockouts Provide a Tool for Examining the Function of Every Gene in an Organism | 569 |
| <i>Summary</i> | 516 | RNA Interference Is a Simple and Rapid Way to Test Gene Function | 571 |
| ANALYZING PROTEINS | 517 | Reporter Genes and <i>In Situ</i> Hybridization Reveal When and Where a Gene Is Expressed | 572 |
| Proteins Can Be Separated by SDS Polyacrylamide-Gel Electrophoresis | 517 | Expression of Individual Genes Can Be Measured Using Quantitative RT-PCR | 573 |
| Specific Proteins Can Be Detected by Blotting with Antibodies | 518 | Microarrays Monitor the Expression of Thousands of Genes at Once | 574 |
| Mass Spectrometry Provides a Highly Sensitive Method for Identifying Unknown Proteins | 519 | Single-Cell Gene Expression Analysis Reveals Biological "Noise" | 575 |
| Two-Dimensional Separation Methods are Especially Powerful | 521 | <i>Summary</i> | 576 |
| Hydrodynamic Measurements Reveal the Size and Shape of a Protein Complex | 522 | <i>Problems</i> | 576 |
| Sets of Interacting Proteins Can Be Identified by Biochemical Methods | 523 | <i>References</i> | 578 |
| Protein-Protein Interactions Can Also Be Identified by a Two-Hybrid Technique in Yeast | 523 | Chapter 9 Visualizing Cells | 579 |
| Combining Data Derived from Different Techniques Produces Reliable Protein-Interaction Maps | 524 | LOOKING AT CELLS IN THE LIGHT MICROSCOPE | 579 |
| Optical Methods Can Monitor Protein Interactions in Real Time | 524 | The Light Microscope Can Resolve Details 0.2 μm Apart | 580 |
| Some Techniques Can Monitor Single Molecules | 526 | Living Cells Are Seen Clearly in a Phase-Contrast or a Differential-Interference-Contrast Microscope | 583 |
| Protein Function Can Be Selectively Disrupted with Small Molecules | 527 | Images Can Be Enhanced and Analyzed by Digital Techniques | 583 |
| Protein Structure Can Be Determined Using X-Ray Diffraction | 527 | Intact Tissues Are Usually Fixed and Sectioned before Microscopy | 585 |
| NMR Can Be Used to Determine Protein Structure in Solution | 529 | Specific Molecules Can Be Located in Cells by Fluorescence Microscopy | 586 |
| Protein Sequence and Structure Provide Clues About Protein Function | 530 | Antibodies Can Be Used to Detect Specific Molecules | 588 |
| <i>Summary</i> | 531 | Imaging of Complex Three-Dimensional Objects Is Possible with the Optical Microscope | 589 |
| ANALYZING AND MANIPULATING DNA | 532 | The Confocal Microscope Produces Optical Sections by Excluding Out-of-Focus Light | 590 |
| Restriction Nucleases Cut Large DNA Molecules into Fragments | 532 | Fluorescent Proteins Can Be Used to Tag Individual Proteins in Living Cells and Organisms | 592 |
| Gel Electrophoresis Separates DNA Molecules of Different Sizes | 534 | Protein Dynamics Can Be Followed in Living Cells | 593 |
| Purified DNA Molecules Can Be Specifically Labeled with Radioisotopes or Chemical Markers <i>in vitro</i> | 535 | Light-Emitting Indicators Can Measure Rapidly Changing Intracellular Ion Concentrations | 596 |
| Nucleic Acid Hybridization Reactions Provide a Sensitive Way of Detecting Specific Nucleotide Sequences | 535 | Several Strategies Are Available by Which Membrane-Impermeant Substances Can Be Introduced into Cells | 597 |
| Northern and Southern Blotting Facilitate Hybridization with Electrophoretically Separated Nucleic Acid Molecules | 538 | Light Can Be Used to Manipulate Microscopic Objects As Well As to Image Them | 598 |
| Genes Can Be Cloned Using DNA Libraries | 540 | Single Molecules Can Be Visualized by Using Total Internal Reflection Fluorescence Microscopy | 599 |
| Two Types of DNA Libraries Serve Different Purposes | 541 | Individual Molecules Can Be Touched and Moved Using Atomic Force Microscopy | 600 |
| cDNA Clones Contain Uninterrupted Coding Sequences | 544 | Molecules Can Be Labeled with Radioisotopes | 600 |
| Genes Can Be Selectively Amplified by PCR | 544 | Radioisotopes Are Used to Trace Molecules in Cells and Organisms | 602 |
| Cells Can Be Used As Factories to Produce Specific Proteins | 546 | <i>Summary</i> | 603 |
| Proteins and Nucleic Acids Can Be Synthesized Directly by Chemical Reactions | 548 | LOOKING AT CELLS AND MOLECULES IN THE ELECTRON MICROSCOPE | 604 |
| DNA Can Be Rapidly Sequenced | 548 | The Electron Microscope Resolves the Fine Structure of the Cell | 604 |
| Nucleotide Sequences Are Used to Predict the Amino Acid Sequences of Proteins | 550 | Biological Specimens Require Special Preparation for the Electron Microscope | 605 |
| The Genomes of Many Organisms Have Been Fully Sequenced | 551 | Specific Macromolecules Can Be Localized by Immunogold Electron Microscopy | 606 |
| <i>Summary</i> | 552 | Images of Surfaces Can Be Obtained by Scanning Electron Microscopy | 607 |
| STUDYING GENE EXPRESSION AND FUNCTION | 553 | Metal Shadowing Allows Surface Features to Be Examined at High Resolution by Transmission Electron Microscopy | 608 |
| Classical Genetics Begins by Disrupting a Cell Process by Random Mutagenesis | 553 | Negative Staining and Cryoelectron Microscopy Both Allow Macromolecules to Be Viewed at High Resolution | 610 |
| Genetic Screens Identify Mutants with Specific Abnormalities | 556 | Multiple Images Can Be Combined to Increase Resolution | 610 |
| Mutations Can Cause Loss or Gain of Protein Function | 557 | Different Views of a Single Object Can Be Combined to Give a Three-Dimensional Reconstruction | 612 |
| Complementation Tests Reveal Whether Two Mutations Are in the Same Gene or Different Genes | 558 | <i>Summary</i> | 612 |
| Genes Can Be Ordered in Pathways by Epistasis Analysis | 558 | <i>Problems</i> | 614 |
| Genes Identified by Mutations Can Be Cloned | 559 | <i>References</i> | 615 |
| Human Genetics Presents Special Problems and Special Opportunities | 560 | Chapter 10 Membrane Structure | 617 |
| Human Genes Are Inherited in Haplotype Blocks, Which Can Aid in the Search for Mutations That Cause Disease | 561 | THE LIPID BILAYER | 617 |
| Complex Traits Are Influenced by Multiple Genes | 563 | Phosphoglycerides, Sphingolipids, and Sterols Are the Major Lipids in Cell Membranes | 618 |
| Reverse Genetics Begins with a Known Gene and Determines Which Cell Processes Require Its Function | 563 | Phospholipids Spontaneously Form Bilayers | 620 |
| Genes Can Be Re-Engineered in Several Ways | 564 | | |
| Engineered Genes Can Be Inserted into the Germ Line of Many Organisms | 565 | | |
| Animals Can Be Genetically Altered | 566 | | |
| Transgenic Plants Are Important for Both Cell Biology and Agriculture | 568 | | |

| | | | |
|--|------------|--|------------|
| The Lipid Bilayer Is a Two-Dimensional Fluid | 621 | Patch-Clamp Recording Indicates That Individual Gated Channels | 680 |
| The Fluidity of a Lipid Bilayer Depends on Its Composition | 622 | Open in an All-or-Nothing Fashion | |
| Despite Their Fluidity, Lipid Bilayers Can Form Domains of Different Compositions | 624 | Voltage-Gated Cation Channels Are Evolutionarily and Structurally Related | 682 |
| Lipid Droplets Are Surrounded by a Phospholipid Monolayer | 625 | Transmitter-Gated Ion Channels Convert Chemical Signals into Electrical Ones at Chemical Synapses | 682 |
| The Asymmetry of the Lipid Bilayer Is Functionally Important | 626 | Chemical Synapses Can Be Excitatory or Inhibitory | 684 |
| Glycolipids Are Found on the Surface of All Plasma Membranes | 628 | The Acetylcholine Receptors at the Neuromuscular Junction Are Transmitter-Gated Cation Channels | 684 |
| <i>Summary</i> | 629 | Transmitter-Gated Ion Channels Are Major Targets for Psychoactive Drugs | 686 |
| MEMBRANE PROTEINS | 629 | Neuromuscular Transmission Involves the Sequential Activation of Five Different Sets of Ion Channels | 687 |
| Membrane Proteins Can Be Associated with the Lipid Bilayer in Various Ways | 629 | Single Neurons Are Complex Computation Devices | 688 |
| Lipid Anchors Control the Membrane Localization of Some Signaling Proteins | 630 | Neuronal Computation Requires a Combination of at Least Three Kinds of K ⁺ Channels | 689 |
| In Most Transmembrane Proteins the Polypeptide Chain Crosses the Lipid Bilayer in an α -Helical Conformation | 631 | Long-Term Potentiation (LTP) in the Mammalian Hippocampus Depends on Ca ²⁺ Entry Through NMDA-Receptor Channels | 691 |
| Transmembrane α Helices Often Interact with One Another | 632 | <i>Summary</i> | 692 |
| Some β Barrels Form Large Transmembrane Channels | 634 | <i>Problems</i> | 693 |
| Many Membrane Proteins Are Glycosylated | 635 | <i>References</i> | 694 |
| Membrane Proteins Can Be Solubilized and Purified in Detergents | 636 | | |
| Bacteriorhodopsin Is a Light-Driven Proton Pump That Traverses the Lipid Bilayer as Seven α Helices | 640 | | |
| Membrane Proteins Often Function as Large Complexes | 642 | Chapter 12 Intracellular Compartments and Protein Sorting | 695 |
| Many Membrane Proteins Diffuse in the Plane of the Membrane | 642 | | |
| Cells Can Confine Proteins and Lipids to Specific Domains Within a Membrane | 645 | THE COMPARTMENTALIZATION OF CELLS | 695 |
| The Cortical Cytoskeleton Gives Membranes Mechanical Strength and Restrict Membrane Protein Diffusion | 646 | All Eucaryotic Cells Have the Same Basic Set of Membrane-Enclosed Organelles | 695 |
| <i>Summary</i> | 648 | Evolutionary Origins Explain the Topological Relationships of Organelles | 697 |
| <i>Problems</i> | 648 | Proteins Can Move Between Compartments in Different Ways | 699 |
| <i>References</i> | 650 | Signal Sequences Direct Proteins to the Correct Cell Address | 701 |
| | | Most Organelles Cannot Be Constructed <i>De Novo</i> : They Require Information in the Organelle Itself | 702 |
| | | <i>Summary</i> | 704 |
| Chapter 11 Membrane Transport of Small Molecules and the Electrical Properties of Membranes | 651 | | |
| | | THE TRANSPORT OF MOLECULES BETWEEN THE NUCLEUS AND THE CYTOSOL | 704 |
| PRINCIPLES OF MEMBRANE TRANSPORT | 651 | Nuclear Pore Complexes Perforate the Nuclear Envelope | 705 |
| Protein-Free Lipid Bilayers Are Highly Impermeable to Ions | 652 | Nuclear Localization Signals Direct Nuclear Proteins to the Nucleus | 705 |
| There Are Two Main Classes of Membrane Transport Proteins: Transporters and Channels | 652 | Nuclear Import Receptors Bind to Both Nuclear Localization Signals and NPC proteins | 707 |
| Active Transport Is Mediated by Transporters Coupled to an Energy Source | 653 | Nuclear Export Works Like Nuclear Import, But in Reverse | 708 |
| <i>Summary</i> | 654 | The Ran GTPase Imposes Directionality on Transport Through NPCs | 708 |
| | | Transport Through NPCs Can Be Regulated by Controlling Access to the Transport Machinery | 709 |
| TRANSPORTERS AND ACTIVE MEMBRANE TRANSPORT | 654 | During Mitosis the Nuclear Envelope Disassembles | 710 |
| Active Transport Can Be Driven by Ion Gradients | 656 | <i>Summary</i> | 712 |
| Transporters in the Plasma Membrane Regulate Cytosolic pH | 657 | | |
| An Asymmetric Distribution of Transporters in Epithelial Cells Underlies the Transcellular Transport of Solutes | 658 | THE TRANSPORT OF PROTEINS INTO MITOCHONDRIA AND CHLOROPLASTS | 713 |
| There Are Three Classes of ATP-Driven Pumps | 659 | Translocation into Mitochondria Depends on Signal Sequences and Protein Translocators | 713 |
| The Ca ²⁺ Pump Is the Best-Understood P-type ATPase | 660 | Mitochondrial Precursor Proteins Are Imported as Unfolded Polypeptide Chains | 715 |
| The Plasma Membrane P-type Na ⁺ -K ⁺ Pump Establishes the Na ⁺ Gradient Across the Plasma Membrane | 661 | ATP Hydrolysis and a Membrane Potential Drive Protein Import Into the Matrix Space | 716 |
| ABC Transporters Constitute the Largest Family of Membrane Transport Proteins | 663 | Bacteria and Mitochondria Use Similar Mechanisms to Insert Porins into their Outer Membrane | 717 |
| <i>Summary</i> | 667 | Transport Into the Inner Mitochondrial Membrane and Intermembrane Space Occurs Via Several Routes | 717 |
| | | Two Signal Sequences Direct Proteins to the Thylakoid Membrane in Chloroplasts | 719 |
| ION CHANNELS AND THE ELECTRICAL PROPERTIES OF MEMBRANES | 667 | <i>Summary</i> | 720 |
| Ion Channels Are Ion-Selective and Fluctuate Between Open and Closed States | 667 | | |
| The Membrane Potential in Animal Cells Depends Mainly on K ⁺ Leak Channels and the K ⁺ Gradient Across the Plasma Membrane | 669 | PEROXISOMES | 721 |
| The Resting Potential Decays Only Slowly When the Na ⁺ -K ⁺ Pump Is Stopped | 669 | Peroxisomes Use Molecular Oxygen and Hydrogen Peroxide to Perform Oxidative Reactions | 721 |
| The Three-Dimensional Structure of a Bacterial K ⁺ Channel Shows How an Ion Channel Can Work | 671 | A Short Signal Sequence Directs the Import of Proteins into Peroxisomes | 722 |
| Aquaporins Are Permeable to Water But Impermeable to Ions | 673 | <i>Summary</i> | 723 |
| The Function of a Neuron Depends on Its Elongated Structure | 675 | | |
| Voltage-Gated Cation Channels Generate Action Potentials in Electrically Excitable Cells | 676 | | |
| Myelination Increases the Speed and Efficiency of Action Potential Propagation in Nerve Cells | 678 | | |

| | | | |
|--|-----|---|-----|
| THE ENDOPLASMIC RETICULUM | 723 | TRANSPORT FROM THE <i>TRANS</i> GOLGI NETWORK TO LYOSOMES | 779 |
| The ER Is Structurally and Functionally Diverse | 724 | Lysosomes Are the Principal Sites of Intracellular Digestion | 779 |
| Signal Sequences Were First Discovered in Proteins Imported into the Rough ER | 726 | Lysosomes Are Heterogeneous | 780 |
| A Signal-Recognition Particle (SRP) Directs ER Signal Sequences to a Specific Receptor in the Rough ER Membrane | 727 | Plant and Fungal Vacuoles Are Remarkably Versatile Lysosomes | 781 |
| The Polypeptide Chain Passes Through an Aqueous Pore in the Translocator | 730 | Multiple Pathways Deliver Materials to Lysosomes | 782 |
| Translocation Across the ER Membrane Does Not Always Require Ongoing Polypeptide Chain Elongation | 731 | A Mannose 6-Phosphate Receptor Recognizes Lysosomal Proteins in the <i>Trans</i> Golgi Network | 783 |
| In Single-Pass Transmembrane Proteins, a Single Internal ER Signal Sequence Remains in the Lipid Bilayer as a Membrane-Spanning α Helix | 732 | The M6P Receptor Shuttles Between Specific Membranes | 784 |
| Combinations of Start-Transfer and Stop-Transfer Signals Determine the Topology of Multipass Transmembrane Proteins | 734 | A Signal Patch in the Hydrolase Polypeptide Chain Provides the Cue for M6P Addition | 785 |
| Translocated Polypeptide Chains Fold and Assemble in the Lumen of the Rough ER | 736 | Defects in the GlcNAc Phosphotransferase Cause a Lysosomal Storage Disease in Humans | 785 |
| Most Proteins Synthesized in the Rough ER Are Glycosylated by the Addition of a Common <i>N</i> -Linked Oligosaccharide | 736 | Some Lysosomes Undergo Exocytosis | 786 |
| Oligosaccharides Are Used as Tags to Mark the State of Protein Folding | 738 | <i>Summary</i> | 786 |
| Improperly Folded Proteins Are Exported from the ER and Degraded in the Cytosol | 739 | TRANSPORT INTO THE CELL FROM THE PLASMA MEMBRANE: ENDOCYTOSIS | 787 |
| Misfolded Proteins in the ER Activate an Unfolded Protein Response | 740 | Specialized Phagocytic Cells Can Ingest Large Particles | 787 |
| Some Membrane Proteins Acquire a Covalently Attached Glycosylphosphatidylinositol (GPI) Anchor | 742 | Pinocytic Vesicles Form from Coated Pits in the Plasma Membrane | 789 |
| The ER Assembles Most Lipid Bilayers | 743 | Not All Pinocytic Vesicles Are Clathrin-Coated | 790 |
| <i>Summary</i> | 745 | Cells Use Receptor-Mediated Endocytosis to Import Selected Extracellular Macromolecules | 791 |
| <i>Problems</i> | 746 | Endocytosed Materials That Are Not Retrieved from Endosomes End Up in Lysosomes | 792 |
| <i>References</i> | 748 | Specific Proteins Are Retrieved from Early Endosomes and Returned to the Plasma Membrane | 793 |
| Chapter 13 Intracellular Vesicular Traffic | 749 | Multivesicular Bodies Form on the Pathway to Late Endosomes | 795 |
| THE MOLECULAR MECHANISMS OF MEMBRANE TRANSPORT AND THE MAINTENANCE OF COMPARTMENTAL DIVERSITY | 750 | Transcytosis Transfers Macromolecules Across Epithelial Cell Sheets | 797 |
| There Are Various Types of Coated Vesicles | 751 | Epithelial Cells Have Two Distinct Early Endosomal Compartments but a Common Late Endosomal Compartment | 798 |
| The Assembly of a Clathrin Coat Drives Vesicle Formation | 754 | <i>Summary</i> | 799 |
| Not All Coats Form Basket-like Structures | 755 | TRANSPORT FROM THE <i>TRANS</i> GOLGI NETWORK TO THE CELL EXTERIOR: EXOCYTOSIS | 799 |
| Phosphoinositides Mark Organelles and Membrane Domains | 757 | Many Proteins and Lipids Seem to Be Carried Automatically from the Golgi Apparatus to the Cell Surface | 800 |
| Cytoplasmic Proteins Regulate the Pinching-Off and Uncoating of Coated Vesicles | 757 | Secretory Vesicles Bud from the <i>Trans</i> Golgi Network | 801 |
| Monomeric GTPases Control Coat Assembly | 758 | Proteins Are Often Proteolytically Processed During the Formation of Secretory Vesicles | 803 |
| Not All Transport Vesicles Are Spherical | 760 | Secretory Vesicles Wait Near the Plasma Membrane Until Signaled to Release Their Contents | 803 |
| Rab Proteins Guide Vesicle Targeting | 760 | Regulated Exocytosis Can Be a Localized Response of the Plasma Membrane and Its Underlying Cytoplasm | 804 |
| SNAREs Mediate Membrane Fusion | 762 | Secretory Vesicle Membrane Components Are Quickly Removed from the Plasma Membrane | 805 |
| Interacting SNAREs Need to Be Pried Apart Before They Can Function Again | 764 | Some Regulated Exocytosis Events Serve to Enlarge the Plasma Membrane | 805 |
| Viral Fusion Proteins and SNAREs May Use Similar Fusion Mechanisms | 764 | Polarized Cells Direct Proteins from the <i>Trans</i> Golgi Network to the Appropriate Domain of the Plasma Membrane | 805 |
| <i>Summary</i> | 766 | Different Strategies Guide Membrane Proteins and Lipids Selectively to the Correct Plasma Membrane Domains | 806 |
| TRANSPORT FROM THE ER THROUGH THE GOLGI APPARATUS | 766 | Synaptic Vesicles Can Form Directly from Endocytic Vesicles | 807 |
| Proteins Leave the ER in COPII-Coated Transport Vesicles | 767 | <i>Summary</i> | 809 |
| Only Proteins That Are Properly Folded and Assembled Can Leave the ER | 767 | <i>Problems</i> | 810 |
| Vesicular Tubular Clusters Mediate Transport from the ER to the Golgi Apparatus | 768 | <i>References</i> | 812 |
| The Retrieval Pathway to the ER Uses Sorting Signals | 769 | Chapter 14 Energy Conversion: Mitochondria and Chloroplasts | 813 |
| Many Proteins Are Selectively Retained in the Compartments in Which They Function | 771 | THE MITOCHONDRION | 815 |
| The Golgi Apparatus Consists of an Ordered Series of Compartments | 771 | The Mitochondrion Contains an Outer Membrane, an Inner Membrane, and Two Internal Compartments | 816 |
| Oligosaccharide Chains Are Processed in the Golgi Apparatus | 773 | The Citric Acid Cycle Generates High-Energy Electrons | 817 |
| Proteoglycans Are Assembled in the Golgi Apparatus | 775 | A Chemiosmotic Process Converts Oxidation Energy into ATP | 817 |
| What Is the Purpose of Glycosylation? | 776 | NADH Transfers its Electrons to Oxygen Through Three Large Respiratory Enzyme Complexes | 819 |
| Transport Through the Golgi Apparatus May Occur by Vesicular Transport or Cisternal Maturation | 777 | As Electrons Move Along the Respiratory Chain, Energy Is Stored as an Electrochemical Proton Gradient Across the Inner Membrane | 820 |
| Golgi Matrix Proteins Help Organize the Stack | 778 | The Proton Gradient Drives ATP Synthesis | 821 |
| <i>Summary</i> | 779 | | |

| | | | |
|---|-----|--|-----|
| The Proton Gradient Drives Coupled Transport Across the Inner Membrane | 822 | Mitochondria and Chloroplasts Have Diverse Genomes | 859 |
| Proton Gradients Produce Most of the Cell's ATP | 822 | Mitochondria and Chloroplasts Probably Both Evolved from Endosymbiotic Bacteria | 859 |
| Mitochondria Maintain a High ATP:ADP Ratio in Cells | 823 | Mitochondria Have a Relaxed Codon Usage and Can Have a Variant Genetic Code | 861 |
| A Large Negative Value of ΔG for ATP Hydrolysis Makes ATP Useful to the Cell | 824 | Animal Mitochondria Contain the Simplest Genetic Systems Known | 862 |
| ATP Synthase Can Function in Reverse to Hydrolyze ATP and Pump H^+ | 826 | Some Organelle Genes Contain Introns | 863 |
| Summary | 827 | The Chloroplast Genome of Higher Plants Contains About 120 Genes | 863 |
| ELECTRON-TRANSPORT CHAINS AND THEIR PROTON PUMPS | 827 | Mitochondrial Genes Are Inherited by a Non-Mendelian Mechanism | 864 |
| Protons Are Unusually Easy to Move | 827 | Organelle Genes Are Maternally Inherited in Many Organisms | 866 |
| The Redox Potential Is a Measure of Electron Affinities | 828 | Petite Mutants in Yeasts Demonstrate the Overwhelming Importance of the Cell Nucleus for Mitochondrial Biogenesis | 866 |
| Electron Transfers Release Large Amounts of Energy | 829 | Mitochondria and Plastids Contain Tissue-Specific Proteins that Are Encoded in the Cell Nucleus | 867 |
| Spectroscopic Methods Identified Many Electron Carriers in the Respiratory Chain | 829 | Mitochondria Import Most of Their Lipids; Chloroplasts Make Most of Theirs | 867 |
| The Respiratory Chain Includes Three Large Enzyme Complexes Embedded in the Inner Membrane | 831 | Mitochondria May Contribute to the Aging of Cells and Organisms | 868 |
| An Iron-Copper Center in Cytochrome Oxidase Catalyzes Efficient O_2 Reduction | 832 | Why Do Mitochondria and Chloroplasts Have Their Own Genetic Systems? | 868 |
| Electron Transfers in the Inner Mitochondrial Membrane Are Mediated by Electron Tunneling during Random Collisions | 834 | Summary | 870 |
| A Large Drop in Redox Potential Across Each of the Three Respiratory Enzyme Complexes Provides the Energy for H^+ Pumping | 835 | THE EVOLUTION OF ELECTRON-TRANSPORT CHAINS | 870 |
| The H^+ Pumping Occurs by Distinct Mechanisms in the Three Major Enzyme Complexes | 835 | The Earliest Cells Probably Used Fermentation to Produce ATP | 870 |
| H^+ Ionophores Uncouple Electron Transport from ATP Synthesis | 836 | Electron-Transport Chains Enabled Anaerobic Bacteria to Use Nonfermentable Molecules as Their Major Source of Energy | 871 |
| Respiratory Control Normally Restrains Electron Flow Through the Chain | 837 | By Providing an Inexhaustible Source of Reducing Power, Photosynthetic Bacteria Overcame a Major Evolutionary Obstacle | 872 |
| Natural Uncouplers Convert the Mitochondria in Brown Fat into Heat-Generating Machines | 838 | The Photosynthetic Electron-Transport Chains of Cyanobacteria Produced Atmospheric Oxygen and Permitted New Life-Forms | 873 |
| The Mitochondrion Plays Many Critical Roles in Cell Metabolism | 838 | Summary | 875 |
| Bacteria Also Exploit Chemiosmotic Mechanisms to Harness Energy | 839 | Problems | 877 |
| Summary | 840 | References | 878 |
| CHLOROPLASTS AND PHOTOSYNTHESIS | 840 | Chapter 15 Mechanisms of Cell Communication | 879 |
| The Chloroplast Is One Member of the Plastid Family of Organelles | 841 | GENERAL PRINCIPLES OF CELL COMMUNICATION | 879 |
| Chloroplasts Resemble Mitochondria But Have an Extra Compartment | 842 | Extracellular Signal Molecules Bind to Specific Receptors | 880 |
| Chloroplasts Capture Energy from Sunlight and Use It to Fix Carbon | 843 | Extracellular Signal Molecules Can Act Over Either Short or Long Distances | 881 |
| Carbon Fixation Is Catalyzed by Ribulose Biphosphate Carboxylase | 844 | Gap Junctions Allow Neighboring Cells to Share Signaling Information | 884 |
| Each CO_2 Molecule That Is Fixed Consumes Three Molecules of ATP and Two Molecules of NADPH | 845 | Each Cell Is Programmed to Respond to Specific Combinations of Extracellular Signal Molecules | 884 |
| Carbon Fixation in Some Plants Is Compartmentalized to Facilitate Growth at Low CO_2 Concentrations | 846 | Different Types of Cells Usually Respond Differently to the Same Extracellular Signal Molecule | 885 |
| Photosynthesis Depends on the Photochemistry of Chlorophyll Molecules | 847 | The Fate of Some Developing Cells Depends on Their Position in Morphogen Gradients | 886 |
| A Photochemical Reaction Center Plus an Antenna Complex Form a Photosystem | 848 | A Cell Can Alter the Concentration of an Intracellular Molecule Quickly Only If the Lifetime of the Molecule Is Short | 886 |
| In a Reaction Center, Light Energy Captured by Chlorophyll Creates a Strong Electron Donor from a Weak One | 849 | Nitric Oxide Gas Signals by Directly Regulating the Activity of Specific Proteins Inside the Target Cell | 887 |
| Noncyclic Photophosphorylation Produces Both NADPH and ATP | 850 | Nuclear Receptors Are Ligand-Modulated Gene Regulatory Proteins | 889 |
| Chloroplasts Can Make ATP by Cyclic Photophosphorylation Without Making NADPH | 853 | The Three Largest Classes of Cell-Surface Receptor Proteins Are Ion-Channel-Coupled, G-Protein-Coupled, and Enzyme-Coupled Receptors | 891 |
| Photosystems I and II Have Related Structures, and Also Resemble Bacterial Photosystems | 853 | Most Activated Cell-Surface Receptors Relay Signals Via Small Molecules and a Network of Intracellular Signaling Proteins | 893 |
| The Proton-Motive Force Is the Same in Mitochondria and Chloroplasts | 853 | Many Intracellular Signaling Proteins Function as Molecular Switches That Are Activated by Phosphorylation or GTP Binding | 895 |
| Carrier Proteins in the Chloroplast Inner Membrane Control Metabolite Exchange with the Cytosol | 854 | Intracellular Signaling Complexes Enhance the Speed, Efficiency, and Specificity of the Response | 897 |
| Chloroplasts Also Perform Other Crucial Biosyntheses | 855 | Modular Interaction Domains Mediate Interactions Between Intracellular Signaling Proteins | 897 |
| Summary | 855 | Cells Can Use Multiple Mechanisms to Respond Abruptly to a Gradually Increasing Concentration of an Extracellular Signal | 899 |
| THE GENETIC SYSTEMS OF MITOCHONDRIA AND PLASTIDS | 855 | Intracellular Signaling Networks Usually Make Use of Feedback Loops | 901 |
| Mitochondria and Chloroplasts Contain Complete Genetic Systems | 856 | Cells Can Adjust Their Sensitivity to a Signal | 902 |
| Organelle Growth and Division Determine the Number of Mitochondria and Plastids in a Cell | 857 | Summary | 903 |

| | | | |
|--|------------|--|-------------|
| SIGNALING THROUGH G-PROTEIN-COUPLED CELL-SURFACE RECEPTORS (GPCRS) AND SMALL INTRACELLULAR MEDIATORS | 904 | Ethylene Blocks the Degradation of Specific Gene Regulatory Proteins in the Nucleus | 957 |
| Trimeric G Proteins Relay Signals from GPCRs | 905 | Regulated Positioning of Auxin Transporters Patterns Plant Growth | 959 |
| Some G Proteins Regulate the Production of Cyclic AMP | 905 | Phytochromes Detect Red Light, and Cryptochromes Detect Blue Light | 960 |
| Cyclic-AMP-Dependent Protein Kinase (PKA) Mediates Most of the Effects of Cyclic AMP | 908 | Summary | 961 |
| Some G Proteins Activate An Inositol Phospholipid Signaling Pathway by Activating Phospholipase C- β | 909 | Problems | 962 |
| Ca ²⁺ Functions as a Ubiquitous Intracellular Mediator | 912 | References | 964 |
| The Frequency of Ca ²⁺ Oscillations Influences a Cell's Response | 912 | | |
| Ca ²⁺ /Calmodulin-Dependent Protein Kinases (CaM-Kinases) Mediate Many of the Responses to Ca ²⁺ Signals in Animal Cells | 914 | Chapter 16 The Cytoskeleton | 965 |
| Some G Proteins Directly Regulate Ion Channels | 916 | THE SELF-ASSEMBLY AND DYNAMIC STRUCTURE OF CYTOSKELETAL FILAMENTS | 965 |
| Smell and Vision Depend on GPCRs That Regulate Cyclic-Nucleotide-Gated Ion Channels | 917 | Cytoskeletal Filaments Are Dynamic and Adaptable | 966 |
| Intracellular Mediators and Enzymatic Cascades Amplify Extracellular Signals | 919 | The Cytoskeleton Can Also Form Stable Structures | 969 |
| GPCR Desensitization Depends on Receptor Phosphorylation | 920 | Each Type of Cytoskeletal Filament Is Constructed from Smaller Protein Subunits | 970 |
| Summary | 921 | Filaments Formed from Multiple Protofilaments Have Advantageous Properties | 971 |
| | | Nucleation Is the Rate-Limiting Step in the Formation of a Cytoskeletal Polymer | 973 |
| SIGNALING THROUGH ENZYME-COUPLED CELL-SURFACE RECEPTORS | 921 | The Tubulin and Actin Subunits Assemble Head-to-Tail to Create Polar Filaments | 973 |
| Activated Receptor Tyrosine Kinases (RTKs) Phosphorylate Themselves | 922 | Microtubules and Actin Filaments Have Two Distinct Ends That Grow at Different Rates | 975 |
| Phosphorylated Tyrosines on RTKs Serve as Docking Sites for Intracellular Signaling Proteins | 923 | Filament Treadmilling and Dynamic Instability Are Consequences of Nucleotide Hydrolysis by Tubulin and Actin | 976 |
| Proteins with SH2 Domains Bind to Phosphorylated Tyrosines | 924 | Treadmilling and Dynamic Instability Aid Rapid Cytoskeletal Rearrangement | 980 |
| Ras Belongs to a Large Superfamily of Monomeric GTPases | 926 | Tubulin and Actin Have Been Highly Conserved During Eucaryotic Evolution | 982 |
| RTKs Activate Ras Via Adaptors and GEFs: Evidence from the Developing <i>Drosophila</i> Eye | 927 | Intermediate Filament Structure Depends on The Lateral Bundling and Twisting of Coiled Coils | 983 |
| Ras Activates a MAP Kinase Signaling Module | 928 | Intermediate Filaments Impart Mechanical Stability to Animal Cells | 985 |
| Scaffold Proteins Help Prevent Cross-Talk Between Parallel MAP Kinase Modules | 930 | Drugs Can Alter Filament Polymerization | 987 |
| Rho Family GTPases Functionally Couple Cell-Surface Receptors to the Cytoskeleton | 931 | Bacterial Cell Organization and Cell Division Depend on Homologs of the Eucaryotic Cytoskeleton | 989 |
| PI 3-Kinase Produces Lipid Docking Sites in the Plasma Membrane | 932 | Summary | 991 |
| The PI-3-Kinase–Akt Signaling Pathway Stimulates Animal Cells to Survive and Grow | 934 | HOW CELLS REGULATE THEIR CYTOSKELETAL FILAMENTS | 992 |
| The Downstream Signaling Pathways Activated By RTKs and GPCRs Overlap | 935 | A Protein Complex Containing γ -Tubulin Nucleates Microtubules | 992 |
| Tyrosine-Kinase-Associated Receptors Depend on Cytoplasmic Tyrosine Kinases | 935 | Microtubules Emanate from the Centrosome in Animal Cells | 992 |
| Cytokine Receptors Activate the JAK–STAT Signaling Pathway, Providing a Fast Track to the Nucleus | 937 | Actin Filaments Are Often Nucleated at the Plasma Membrane | 996 |
| Protein Tyrosine Phosphatases Reverse Tyrosine Phosphorylations | 938 | The Mechanism of Nucleation Influences Large-Scale Filament Organization | 998 |
| Signal Proteins of the TGF β Superfamily Act Through Receptor Serine/Threonine Kinases and Smads | 939 | Proteins That Bind to the Free Subunits Modify Filament Elongation | 999 |
| Serine/Threonine and Tyrosine Protein Kinases Are Structurally Related | 941 | Severing Proteins Regulate the Length and Kinetic Behavior of Actin Filaments and Microtubules | 1000 |
| Bacterial Chemotaxis Depends on a Two-Component Signaling Pathway Activated by Histidine-Kinase-Associated Receptors | 941 | Proteins That Bind Along the Sides of Filaments Can Either Stabilize or Destabilize Them | 1001 |
| Receptor Methylation Is Responsible for Adaptation in Bacterial Chemotaxis | 943 | Proteins That Interact with Filament Ends Can Dramatically Change Filament Dynamics | 1002 |
| Summary | 944 | Different Kinds of Proteins Alter the Properties of Rapidly Growing Microtubule Ends | 1003 |
| | | Filaments Are Organized into Higher-Order Structures in Cells | 1005 |
| SIGNALING PATHWAYS DEPENDENT ON REGULATED PROTEOLYSIS OF LATENT GENE REGULATORY PROTEINS | 946 | Intermediate Filaments Are Cross-Linked and Bundled Into Strong Arrays | 1005 |
| The Receptor Protein Notch Is a Latent Gene Regulatory Protein | 946 | Cross-Linking Proteins with Distinct Properties Organize Different Assemblies of Actin Filaments | 1006 |
| Wnt Proteins Bind to Frizzled Receptors and Inhibit the Degradation of β -Catenin | 948 | Filamin and Spectrin Form Actin Filament Webs | 1008 |
| Hedgehog Proteins Bind to Patched Relieving Its Inhibition of Smoothened | 950 | Cytoskeletal Elements Make Many Attachments to Membrane | 1009 |
| Many Stressful and Inflammatory Stimuli Act Through an NF κ B-Dependent Signaling Pathway | 952 | Summary | 1010 |
| Summary | 954 | MOLECULAR MOTORS | 1010 |
| | | Actin-Based Motor Proteins Are Members of the Myosin Superfamily | 1011 |
| SIGNALING IN PLANTS | 955 | There Are Two Types of Microtubule Motor Proteins: Kinesins and Dyneins | 1014 |
| Multicellularity and Cell Communication Evolved Independently in Plants and Animals | 955 | The Structural Similarity of Myosin and Kinesin Indicates a Common Evolutionary Origin | 1015 |
| Receptor Serine/Threonine Kinases Are the Largest Class of Cell-Surface Receptors in Plants | 956 | Motor Proteins Generate Force by Coupling ATP Hydrolysis to Conformational Changes | 1016 |

| | | | |
|--|------|---|------|
| Motor Protein Kinetics Are Adapted to Cell Functions | 1020 | Microtubule-Dependent Motor Proteins Govern Spindle Assembly and Function | 1077 |
| Motor Proteins Mediate the Intracellular Transport of Membrane-Enclosed Organelles | 1021 | Two Mechanisms Collaborate in the Assembly of a Bipolar Mitotic Spindle | 1077 |
| The Cytoskeleton Localizes Specific RNA Molecules | 1022 | Centrosome Duplication Occurs Early in the Cell Cycle | 1078 |
| Cells Regulate Motor Protein Function | 1023 | M-Cdk Initiates Spindle Assembly in Prophase | 1078 |
| <i>Summary</i> | 1025 | The Completion of Spindle Assembly in Animal Cells Requires Nuclear Envelope Breakdown | 1079 |
| THE CYTOSKELETON AND CELL BEHAVIOR | 1025 | Microtubule Instability Increases Greatly in Mitosis | 1080 |
| Sliding of Myosin II and Actin Filaments Causes Muscles to Contract | 1026 | Mitotic Chromosomes Promote Bipolar Spindle Assembly | 1081 |
| A Sudden Rise in Cytosolic Ca ²⁺ Concentration Initiates Muscle Contraction | 1028 | Kinetochores Attach Sister Chromatids to the Spindle | 1082 |
| Heart Muscle Is a Precisely Engineered Machine | 1031 | Bi-Orientation Is Achieved by Trial and Error | 1083 |
| Cilia and Flagella Are Motile Structures Built from Microtubules and Dyneins | 1031 | Multiple Forces Move Chromosomes on the Spindle | 1085 |
| Construction of the Mitotic Spindle Requires Microtubule Dynamics and the Interactions of Many Motor Proteins | 1034 | The APC/C Triggers Sister-Chromatid Separation and the Completion of Mitosis | 1087 |
| Many Cells Can Crawl Across A Solid Substratum | 1036 | Unattached Chromosomes Block Sister-Chromatid Separation: The Spindle Assembly Checkpoint | 1088 |
| Actin Polymerization Drives Plasma Membrane Protrusion | 1037 | Chromosomes Segregate in Anaphase A and B | 1089 |
| Cell Adhesion and Traction Allow Cells to Pull Themselves Forward | 1040 | Segregated Chromosomes Are Packaged in Daughter Nuclei at Telophase | 1090 |
| Members of the Rho Protein Family Cause Major Rearrangements of the Actin Cytoskeleton | 1041 | Meiosis Is a Special Form of Nuclear Division Involved in Sexual Reproduction | 1090 |
| Extracellular Signals Can Activate the Three Rho Protein Family Members | 1043 | <i>Summary</i> | 1092 |
| External Signals Can Dictate the Direction of Cell Migration | 1045 | CYTOKINESIS | 1092 |
| Communication Between the Microtubule and Actin Cytoskeletons Coordinates Whole-Cell Polarization and Locomotion | 1046 | Actin and Myosin II in the Contractile Ring Generate the Force for Cytokinesis | 1093 |
| The Complex Morphological Specialization of Neurons Depends on the Cytoskeleton | 1047 | Local Activation of RhoA Triggers Assembly and Contraction of the Contractile Ring | 1094 |
| <i>Summary</i> | 1050 | The Microtubules of the Mitotic Spindle Determine the Plane of Animal Cell Division | 1095 |
| <i>Problems</i> | 1050 | The Phragmoplast Guides Cytokinesis in Higher Plants | 1097 |
| <i>References</i> | 1052 | Membrane-Enclosed Organelles Must Be Distributed to Daughter Cells During Cytokinesis | 1098 |
| Chapter 17 The Cell Cycle | 1053 | Some Cells Reposition Their Spindle to Divide Asymmetrically | 1099 |
| OVERVIEW OF THE CELL CYCLE | 1054 | Mitosis Can Occur Without Cytokinesis | 1099 |
| The Eucaryotic Cell Cycle Is Divided into Four Phases | 1054 | The G ₁ Phase Is a Stable State of Cdk Inactivity | 1100 |
| Cell-Cycle Control Is Similar in All Eucaryotes | 1056 | <i>Summary</i> | 1101 |
| Cell-Cycle Control Can Be Dissected Genetically by Analysis of Yeast Mutants | 1056 | CONTROL OF CELL DIVISION AND CELL GROWTH | 1101 |
| Cell-Cycle Control Can Be Analyzed Biochemically in Animal Embryos | 1057 | Mitogens Stimulate Cell Division | 1102 |
| Cell-Cycle Control Can Be Studied in Cultured Mammalian Cells | 1059 | Cells Can Delay Division by Entering a Specialized Nondividing State | 1103 |
| Cell-Cycle Progression Can Be Studied in Various Ways | 1059 | Mitogens Stimulate G ₁ -Cdk and G ₁ /S-Cdk Activities | 1103 |
| <i>Summary</i> | 1060 | DNA Damage Blocks Cell Division: The DNA Damage Response | 1105 |
| THE CELL-CYCLE CONTROL SYSTEM | 1060 | Many Human Cells Have a Built-In Limitation on the Number of Times They Can Divide | 1007 |
| The Cell-Cycle Control System Triggers the Major Events of the Cell Cycle | 1060 | Abnormal Proliferation Signals Cause Cell-Cycle Arrest or Apoptosis, Except in Cancer Cells | 1107 |
| The Cell-Cycle Control System Depends on Cyclically Activated Cyclin-Dependent Protein Kinases (Cdks) | 1062 | Organism and Organ Growth Depend on Cell Growth | 1108 |
| Inhibitory Phosphorylation and Cdk Inhibitory Proteins (CKIs) Can Suppress Cdk Activity | 1063 | Proliferating Cells Usually Coordinate Their Growth and Division | 1108 |
| The Cell-Cycle Control System Depends on Cyclical Proteolysis | 1064 | Neighboring Cells Compete for Extracellular Signal Proteins | 1110 |
| Cell-Cycle Control Also Depends on Transcriptional Regulation | 1065 | Animals Control Total Cell Mass by Unknown Mechanisms | 1111 |
| The Cell-Cycle Control System Functions as a Network of Biochemical Switches | 1065 | <i>Summary</i> | 1112 |
| <i>Summary</i> | 1067 | <i>Problems</i> | 1112 |
| S PHASE | 1067 | <i>References</i> | 1113 |
| S-Cdk Initiates DNA Replication Once Per Cycle | 1067 | Chapter 18 Apoptosis | 1115 |
| Chromosome Duplication Requires Duplication of Chromatin Structure | 1069 | Programmed Cell Death Eliminates Unwanted Cells | 1115 |
| Cohesins Help Hold Sister Chromatids Together | 1070 | Apoptotic Cells Are Biochemically Recognizable | 1117 |
| <i>Summary</i> | 1071 | Apoptosis Depends on an Intracellular Proteolytic Cascade That Is Mediated by Caspases | 1118 |
| MITOSIS | 1071 | Cell-Surface Death Receptors Activate the Extrinsic Pathway of Apoptosis | 1120 |
| M-Cdk Drives Entry Into Mitosis | 1071 | The Intrinsic Pathway of Apoptosis Depends on Mitochondria | 1121 |
| Dephosphorylation Activates M-Cdk at the Onset of Mitosis | 1074 | Bcl2 Proteins Regulate the Intrinsic Pathway of Apoptosis | 1121 |
| Condensin Helps Configure Duplicated Chromosomes for Separation | 1075 | IAPs Inhibit Caspases | 1124 |
| The Mitotic Spindle Is a Microtubule-Based Machine | 1075 | Extracellular Survival Factors Inhibit Apoptosis in Various Ways | 1126 |
| | | Either Excessive or Insufficient Apoptosis Can Contribute to Disease | 1127 |
| | | <i>Summary</i> | 1128 |
| | | <i>Problems</i> | 1128 |
| | | <i>References</i> | 1129 |

Chapter 19 Cell Junctions, Cell Adhesion, and the Extracellular Matrix

| | |
|--|-------------|
| CADHERINS AND CELL–CELL ADHESION | 1133 |
| Cadherins Mediate Ca ²⁺ -Dependent Cell–Cell Adhesion in All Animals | 1135 |
| The Cadherin Superfamily in Vertebrates Includes Hundreds of Different Proteins, Including Many with Signaling Functions | 1136 |
| Cadherins Mediate Homophilic Adhesion | 1137 |
| Selective Cell–Cell Adhesion Enables Dissociated Vertebrate Cells to Reassemble into Organized Tissues | 1139 |
| Cadherins Control the Selective Assortment of Cells | 1140 |
| Twist Regulates Epithelial–Mesenchymal Transitions | 1141 |
| Catenins Link Classical Cadherins to the Actin Cytoskeleton | 1142 |
| Adherens Junctions Coordinate the Actin-Based Motility of Adjacent Cells | 1142 |
| Desmosome Junctions Give Epithelia Mechanical Strength | 1143 |
| Cell–Cell Junctions Send Signals into the Cell Interior | 1145 |
| Selectins Mediate Transient Cell–Cell Adhesions in the Bloodstream | 1145 |
| Members of the Immunoglobulin Superfamily of Proteins Mediate Ca ²⁺ -Independent Cell–Cell Adhesion | 1146 |
| Many Types of Cell Adhesion Molecules Act in Parallel to Create a Synapse | 1147 |
| Scaffold Proteins Organize Junctional Complexes | 1148 |
| <i>Summary</i> | 1149 |
| TIGHT JUNCTIONS AND THE ORGANIZATION OF EPITHELIA | 1150 |
| Tight Junctions Form a Seal Between Cells and a Fence Between Membrane Domains | 1150 |
| Scaffold Proteins in Junctional Complexes Play a Key Part in the Control of Cell Proliferation | 1153 |
| Cell–Cell Junctions and the Basal Lamina Govern Apico-Basal Polarity in Epithelia | 1155 |
| A Separate Signaling System Controls Planar Cell Polarity | 1157 |
| <i>Summary</i> | 1158 |
| PASSAGEWAYS FROM CELL TO CELL: GAP JUNCTIONS AND PLASMODESMATA | 1158 |
| Gap Junctions Couple Cells Both Electrically and Metabolically | 1158 |
| A Gap-Junction Connexon Is Made Up of Six Transmembrane Connexin Subunits | 1159 |
| Gap Junctions Have Diverse Functions | 1161 |
| Cells Can Regulate the Permeability of Their Gap Junctions | 1161 |
| In Plants, Plasmodesmata Perform Many of the Same Functions as Gap Junctions | 1162 |
| <i>Summary</i> | 1163 |
| THE BASAL LAMINA | 1164 |
| Basal Laminae Underlie All Epithelia and Surround Some Nonepithelial Cell Types | 1164 |
| Laminin Is a Primary Component of the Basal Lamina | 1165 |
| Type IV Collagen Gives the Basal Lamina Tensile Strength | 1166 |
| Basal Laminae Have Diverse Functions | 1167 |
| <i>Summary</i> | 1169 |
| INTEGRINS AND CELL–MATRIX ADHESION | 1169 |
| Integrins Are Transmembrane Heterodimers That Link to the Cytoskeleton | 1170 |
| Integrins Can Switch Between an Active and an Inactive Conformation | 1170 |
| Integrin Defects Are Responsible for Many Different Genetic Diseases | 1172 |
| Integrins Cluster to Form Strong Adhesions | 1174 |
| Extracellular Matrix Attachments Act Through Integrins to Control Cell Proliferation and Survival | 1175 |
| Integrins Recruit Intracellular Signaling Proteins at Sites of Cell–Substratum Adhesion | 1176 |
| Integrins Can Produce Localized Intracellular Effects | 1177 |
| <i>Summary</i> | 1178 |

| | |
|---|-------------|
| THE EXTRACELLULAR MATRIX OF ANIMAL CONNECTIVE TISSUES | 1178 |
| The Extracellular Matrix Is Made and Oriented by the Cells Within It | 1179 |
| Glycosaminoglycan (GAG) Chains Occupy Large Amounts of Space and Form Hydrated Gels | 1179 |
| Hyaluronan Acts as a Space Filler and a Facilitator of Cell Migration During Tissue Morphogenesis and Repair | 1180 |
| Proteoglycans Are Composed of GAG Chains Covalently Linked to a Core Protein | 1181 |
| Proteoglycans Can Regulate the Activities of Secreted Proteins | 1182 |
| Cell-Surface Proteoglycans Act as Co-Receptors | 1183 |
| Collagens Are the Major Proteins of the Extracellular Matrix | 1184 |
| Collagen Chains Undergo a Series of Post-Translational Modifications | 1186 |
| Propeptides Are Clipped Off Procollagen After Its Secretion to Allow Assembly of Fibrils | 1187 |
| Secreted Fibril-Associated Collagens Help Organize the Fibrils | 1187 |
| Cells Help Organize the Collagen Fibrils They Secrete by Exerting Tension on the Matrix | 1189 |
| Elastin Gives Tissues Their Elasticity | 1189 |
| Fibronectin Is an Extracellular Protein That Helps Cells Attach to the Matrix | 1191 |
| Tension Exerted by Cells Regulates Assembly of Fibronectin Fibrils | 1191 |
| Fibronectin Binds to Integrins Through an RGD Motif | 1193 |
| Cells Have to Be Able to Degrade Matrix, as Well as Make it | 1193 |
| Matrix Degradation Is Localized to the Vicinity of Cells | 1194 |
| <i>Summary</i> | 1195 |
| THE PLANT CELL WALL | 1195 |
| The Composition of the Cell Wall Depends on the Cell Type | 1195 |
| The Tensile Strength of the Cell Wall Allows Plant Cells to Develop Turgor Pressure | 1197 |
| The Primary Cell Wall Is Built from Cellulose Microfibrils Interwoven with a Network of Pectic Polysaccharides | 1197 |
| Oriented Cell-Wall Deposition Controls Plant Cell Growth | 1199 |
| Microtubules Orient Cell-Wall Deposition | 1200 |
| <i>Summary</i> | 1202 |
| <i>Problems</i> | 1202 |
| <i>References</i> | 1204 |
| Chapter 20 Cancer | 1205 |
| CANCER AS A MICROEVOLUTIONARY PROCESS | 1205 |
| Cancer Cells Reproduce Without Restraint and Colonize Other Tissues | 1206 |
| Most Cancers Derive from a Single Abnormal Cell | 1207 |
| Cancer Cells Contain Somatic Mutations | 1208 |
| A Single Mutation Is Not Enough to Cause Cancer | 1209 |
| Cancers Develop Gradually from Increasingly Aberrant Cells | 1210 |
| Cervical Cancers Are Prevented by Early Detection | 1211 |
| Tumor Progression Involves Successive Rounds of Random Inherited Change Followed by Natural Selection | 1212 |
| The Epigenetic Changes That Accumulate in Cancer Cells Involve Inherited Chromatin Structures and DNA Methylation | 1213 |
| Human Cancer Cells Are Genetically Unstable | 1214 |
| Cancerous Growth Often Depends on Defective Control of Cell Death, Cell Differentiation, or Both | 1215 |
| Cancer Cells Are Usually Altered in Their Responses to DNA Damage and Other Forms of Stress | 1216 |
| Human Cancer Cells Escape a Built-In Limit to Cell Proliferation | 1217 |
| A Small Population of Cancer Stem Cells Maintains Many Tumors | 1217 |
| How Do Cancer Stem Cells Arise? | 1218 |
| To Metastasize, Malignant Cancer Cells Must Survive and Proliferate in a Foreign Environment | 1220 |
| Tumors Induce Angiogenesis | 1220 |
| The Tumor Microenvironment Influences Cancer Development | 1222 |
| Many Properties Typically Contribute to Cancerous Growth | 1223 |
| <i>Summary</i> | 1223 |

| | | | |
|---|------|--|------|
| THE PREVENTABLE CAUSES OF CANCER | 1224 | There Is Still Much More to Do | 1264 |
| Many, But Not All, Cancer-Causing Agents Damage DNA | 1225 | <i>Summary</i> | 1265 |
| Tumor Initiators Damage DNA; Tumor Promoters Do Not | 1226 | <i>Problems</i> | 1265 |
| Viruses and Other Infections Contribute to a Significant Proportion of Human Cancers | 1227 | <i>References</i> | 1267 |
| Identification of Carcinogens Reveals Ways to Avoid Cancer | 1229 | | |
| <i>Summary</i> | 1230 | | |
| FINDING THE CANCER-CRITICAL GENES | 1230 | Chapter 21 Sexual Reproduction: Meiosis, Germ Cells, and Fertilization | 1269 |
| The Identification of Gain-of-Function and Loss-of-Function Mutations Requires Different Methods | 1231 | OVERVIEW OF SEXUAL REPRODUCTION | 1269 |
| Retroviruses Can Act as Vectors for Oncogenes That Alter Cell Behavior | 1232 | The Haploid Phase in Higher Eucaryotes Is Brief | 1269 |
| Different Searches for Oncogenes Have Converged on the Same Gene—Ras | 1233 | Meiosis Creates Genetic Diversity | 1271 |
| Studies of Rare Hereditary Cancer Syndromes First Identified Tumor Suppressor Genes | 1234 | Sexual Reproduction Gives Organisms a Competitive Advantage | 1271 |
| Tumor Suppressor Genes Can Also Be Identified from Studies of Tumors | 1235 | <i>Summary</i> | 1272 |
| Both Genetic and Epigenetic Mechanisms Can Inactivate Tumor Suppressor Genes | 1235 | MEIOSIS | 1272 |
| Genes Mutated in Cancer Can Be Made Overactive in Many Ways | 1237 | Gametes Are Produced by Two Meiotic Cell Divisions | 1272 |
| The Hunt for Cancer-Critical Genes Continues | 1239 | Duplicated Homologs (and Sex Chromosomes) Pair During Early Prophase I | 1274 |
| <i>Summary</i> | 1240 | Homolog Pairing Culminates in the Formation of a Synaptonemal Complex | 1275 |
| THE MOLECULAR BASIS OF CANCER-CELL BEHAVIOR | 1240 | Homolog Segregation Depends on Meiosis-Specific, Kinetochore-Associated Proteins | 1276 |
| Studies of Both Developing Embryos and Genetically Engineered Mice Have Helped to Uncover the Function of Cancer-Critical Genes | 1241 | Meiosis Frequently Goes Wrong | 1278 |
| Many Cancer-Critical Genes Regulate Cell Proliferation | 1242 | Crossing-Over Enhances Genetic Reassortment | 1279 |
| Distinct Pathways May Mediate the Disregulation of Cell-Cycle Progression and the Disregulation of Cell Growth in Cancer Cells | 1244 | Crossing-Over Is Highly Regulated | 1280 |
| Mutations in Genes That Regulate Apoptosis Allow Cancer Cells to Survive When They Should Not | 1245 | Meiosis Is Regulated Differently in Male and Female Mammals | 1280 |
| Mutations in the p53 Gene Allow Many Cancer Cells to Survive and Proliferate Despite DNA Damage | 1246 | <i>Summary</i> | 1281 |
| DNA Tumor Viruses Block the Action of Key Tumor Suppressor Proteins | 1247 | PRIMORDIAL GERM CELLS AND SEX DETERMINATION IN MAMMALS | 1282 |
| The Changes in Tumor Cells That Lead to Metastasis Are Still Largely a Mystery | 1249 | Signals from Neighbors Specify PGCs in Mammalian Embryos | 1282 |
| Colorectal Cancers Evolve Slowly Via a Succession of Visible Changes | 1250 | PGCs Migrate into the Developing Gonads | 1283 |
| A Few Key Genetic Lesions Are Common to a Large Fraction of Colorectal Cancers | 1251 | The Sry Gene Directs the Developing Mammalian Gonad to Become a Testis | 1283 |
| Some Colorectal Cancers Have Defects in DNA Mismatch Repair | 1254 | Many Aspects of Sexual Reproduction Vary Greatly between Animal Species | 1285 |
| The Steps of Tumor Progression Can Often Be Correlated with Specific Mutations | 1254 | <i>Summary</i> | 1286 |
| Each Case of Cancer Is Characterized by Its Own Array of Genetic Lesions | 1256 | EGGS | 1287 |
| <i>Summary</i> | 1256 | An Egg Is Highly Specialized for Independent Development | 1287 |
| CANCER TREATMENT: PRESENT AND FUTURE | 1256 | Eggs Develop in Stages | 1288 |
| The Search for Cancer Cures Is Difficult but Not Hopeless | 1257 | Oocytes Use Special Mechanisms to Grow to Their Large Size | 1290 |
| Traditional Therapies Exploit the Genetic Instability and Loss of Cell-Cycle Checkpoint Responses in Cancer Cells | 1257 | Most Human Oocytes Die Without Maturing | 1291 |
| New Drugs Can Exploit the Specific Cause of a Tumor's Genetic Instability | 1257 | <i>Summary</i> | 1292 |
| Genetic Instability Helps Cancers Become Progressively More Resistant to Therapies | 1259 | SPERM | 1292 |
| New Therapies Are Emerging from Our Knowledge of Cancer Biology | 1260 | Sperm Are Highly Adapted for Delivering Their DNA to an Egg | 1292 |
| Small Molecules Can Be Designed to Inhibit Specific Oncogenic Proteins | 1260 | Sperm Are Produced Continuously in the Mammalian Testis | 1293 |
| Tumor Blood Vessels Are Logical Targets for Cancer Therapy | 1262 | Sperm Develop as a Syncytium | 1294 |
| Many Cancers May Be Treatable by Enhancing the Immune Response Against a Specific Tumor | 1262 | <i>Summary</i> | 1296 |
| Treating Patients with Several Drugs Simultaneously Has Potential Advantages for Cancer Therapy | 1263 | FERTILIZATION | 1297 |
| Gene Expression Profiling Can Help Classify Cancers into Clinically Meaningful Subgroups | 1264 | Ejaculated Sperm Become Capacitated in the Female Genital Tract | 1297 |
| | | Capacitated Sperm Bind to the Zona Pellucida and Undergo an Acrosome Reaction | 1298 |
| | | The Mechanism of Sperm–Egg Fusion Is Still Unknown | 1298 |
| | | Sperm Fusion Activates the Egg by Increasing Ca ²⁺ in the Cytosol | 1299 |
| | | The Cortical Reaction Helps Ensure That Only One Sperm Fertilizes the Egg | 1300 |
| | | The Sperm Provides Centrioles as Well as Its Genome to the Zygote | 1301 |
| | | IVF and ICSI Have Revolutionized the Treatment of Human Infertility | 1301 |
| | | <i>Summary</i> | 1303 |
| | | <i>References</i> | 1304 |
| | | Chapter 22 Development of Multicellular Organisms | 1305 |
| | | UNIVERSAL MECHANISMS OF ANIMAL DEVELOPMENT | 1305 |
| | | Animals Share Some Basic Anatomical Features | 1307 |

| | | | |
|--|------|---|------|
| Multicellular Animals Are Enriched in Proteins Mediating Cell Interactions and Gene Regulation | 1308 | Egg-Polarity, Gap, and Pair-Rule Genes Create a Transient Pattern That Is Remembered by Other Genes | 1340 |
| Regulatory DNA Defines the Program of Development | 1309 | Summary | 1341 |
| Manipulation of the Embryo Reveals the Interactions Between Its Cells | 1310 | HOMEOTIC SELECTOR GENES AND THE PATTERNING OF THE ANTEROPOSTERIOR AXIS | 1341 |
| Studies of Mutant Animals Identify the Genes That Control Developmental Processes | 1311 | The Hox Code Specifies Anterior-Posterior Differences | 1342 |
| A Cell Makes Developmental Decisions Long Before It Shows a Visible Change | 1311 | Homeotic Selector Genes Code for DNA-Binding Proteins That Interact with Other Gene Regulatory Proteins | 1342 |
| Cells Have Remembered Positional Values That Reflect Their Location in the Body | 1312 | The Homeotic Selector Genes Are Expressed Sequentially According to Their Order in the Hox Complex | 1343 |
| Inductive Signals Can Create Orderly Differences Between Initially Identical Cells | 1313 | The Hox Complex Carries a Permanent Record of Positional Information | 1344 |
| Sister Cells Can Be Born Different by an Asymmetric Cell Division | 1313 | The Anteroposterior Axis Is Controlled by <i>Hox</i> Selector Genes in Vertebrates Also | 1344 |
| Positive Feedback Can Create Asymmetry Where There Was None Before | 1314 | Summary | 1347 |
| Positive Feedback Generates Patterns, Creates All-or-None Outcomes, and Provides Memory | 1315 | ORGANOGENESIS AND THE PATTERNING OF APPENDAGES | 1347 |
| A Small Set of Signaling Pathways, Used Repeatedly, Controls Developmental Patterning | 1316 | Conditional and Induced Somatic Mutations Make it Possible to Analyze Gene Functions Late in Development | 1348 |
| Morphogens Are Long-Range Inducers That Exert Graded Effects | 1316 | Body Parts of the Adult Fly Develop From Imaginal Discs | 1349 |
| Extracellular Inhibitors of Signal Molecules Shape the Response to the Inducer | 1317 | Homeotic Selector Genes Are Essential for the Memory of Positional Information in Imaginal Disc Cells | 1351 |
| Developmental Signals Can Spread Through Tissue in Several Different Ways | 1318 | Specific Regulatory Genes Define the Cells That Will Form an Appendage | 1351 |
| Programs That Are Intrinsic to a Cell Often Define the Time-Course of its Development | 1319 | The Insect Wing Disc Is Divided into Compartments | 1352 |
| Initial Patterns Are Established in Small Fields of Cells and Refined by Sequential Induction as the Embryo Grows | 1319 | Four Familiar Signaling Pathways Combine to Pattern the Wing Disc: Wingless, Hedgehog, Dpp, and Notch | 1353 |
| Summary | 1320 | The Size of Each Compartment Is Regulated by Interactions Among Its Cells | 1353 |
| CAENORHABDITIS ELEGANS: DEVELOPMENT FROM THE PERSPECTIVE OF THE INDIVIDUAL CELL | 1321 | Similar Mechanisms Pattern the Limbs of Vertebrates | 1355 |
| <i>Caenorhabditis elegans</i> Is Anatomically Simple | 1321 | Localized Expression of Specific Classes of Gene Regulatory Proteins Foreshadows Cell Differentiation | 1356 |
| Cell Fates in the Developing Nematode Are Almost Perfectly Predictable | 1322 | Lateral Inhibition Singles Out Sensory Mother Cells Within Proneural Clusters | 1357 |
| Products of Maternal-Effect Genes Organize the Asymmetric Division of the Egg | 1323 | Lateral Inhibition Drives the Progeny of the Sensory Mother Cell Toward Different Final Fates | 1357 |
| Progressively More Complex Patterns Are Created by Cell-Cell Interactions | 1324 | Planar Polarity of Asymmetric Divisions Is Controlled by Signaling via the Receptor Frizzled | 1358 |
| Microsurgery and Genetics Reveal the Logic of Developmental Control; Gene Cloning and Sequencing Reveal Its Molecular Mechanisms | 1325 | Asymmetric Stem-Cell Divisions Generate Additional Neurons in the Central Nervous System | 1359 |
| Cells Change Over Time in Their Responsiveness to Developmental Signals | 1325 | Asymmetric Neuroblast Divisions Segregate an Inhibitor of Cell Division into Just One of the Daughter Cells | 1361 |
| Heterochronic Genes Control the Timing of Development | 1326 | Notch Signaling Regulates the Fine-Grained Pattern of Differentiated Cell Types in Many Different Tissues | 1362 |
| Cells Do Not Count Cell Divisions in Timing Their Internal Programs | 1327 | Some Key Regulatory Genes Define a Cell Type; Others Can Activate the Program for Creation of an Entire Organ | 1362 |
| Selected Cells Die by Apoptosis as Part of the Program of Development | 1327 | Summary | 1363 |
| Summary | 1328 | CELL MOVEMENTS AND THE SHAPING OF THE VERTEBRATE BODY | 1363 |
| DROSOPHILA AND THE MOLECULAR GENETICS OF PATTERN FORMATION: GENESIS OF THE BODY PLAN | 1328 | The Polarity of the Amphibian Embryo Depends on the Polarity of the Egg | 1364 |
| The Insect Body Is Constructed as a Series of Segmental Units | 1329 | Cleavage Produces Many Cells from One | 1365 |
| <i>Drosophila</i> Begins Its Development as a Syncytium | 1330 | Gastrulation Transforms a Hollow Ball of Cells into a Three-Layered Structure with a Primitive Gut | 1365 |
| Genetic Screens Define Groups of Genes Required for Specific Aspects of Early Patterning | 1332 | The Movements of Gastrulation Are Precisely Predictable | 1366 |
| Interactions of the Oocyte With Its Surroundings Define the Axes of the Embryo: the Role of the Egg-Polarity Genes | 1333 | Chemical Signals Trigger the Mechanical Processes | 1367 |
| The Dorsoventral Signaling Genes Create a Gradient of a Nuclear Gene Regulatory Protein | 1334 | Active Changes of Cell Packing Provide a Driving Force for Gastrulation | 1368 |
| Dpp and Sog Set Up a Secondary Morphogen Gradient to Refine the Pattern of the Dorsal Part of the Embryo | 1336 | Changing Patterns of Cell Adhesion Molecules Force Cells Into New Arrangements | 1369 |
| The Insect Dorsoventral Axis Corresponds to the Vertebrate Ventrordorsal Axis | 1336 | The Notochord Elongates, While the Neural Plate Rolls Up to Form the Neural Tube | 1370 |
| Three Classes of Segmentation Genes Refine the Anterior-Posterior Maternal Pattern and Subdivide the Embryo | 1336 | A Gene-Expression Oscillator Controls Segmentation of the Mesoderm Into Somites | 1371 |
| The Localized Expression of Segmentation Genes Is Regulated by a Hierarchy of Positional Signals | 1337 | Delayed Negative Feedback May Generate the Oscillations of the Segmentation Clock | 1373 |
| The Modular Nature of Regulatory DNA Allows Genes to Have Multiple Independently Controlled Functions | 1339 | Embryonic Tissues Are Invaded in a Strictly Controlled Fashion by Migratory Cells | 1373 |
| | | The Distribution of Migrant Cells Depends on Survival Factors as Well as Guidance Cues | 1375 |

| | | | |
|---|------|---|------|
| Left-Right Asymmetry of the Vertebrate Body Derives From Molecular Asymmetry in the Early Embryo | 1376 | The Basal Layer Contains Both Stem Cells and Transit Amplifying Cells | 1422 |
| <i>Summary</i> | 1377 | Transit amplifying Divisions Are Part of the Strategy of Growth Control | 1423 |
| THE MOUSE | 1378 | Stem Cells of Some Tissues Selectively Retain Original DNA Strands | 1424 |
| Mammalian Development Begins With a Specialized Preamble | 1378 | The Rate of Stem-Cell Division Can Increase Dramatically When New Cells Are Needed Urgently | 1425 |
| The Early Mammalian Embryo Is Highly Regulative | 1380 | Many Interacting Signals Govern Epidermal Renewal | 1426 |
| Totipotent Embryonic Stem Cells Can Be Obtained From a Mammalian Embryo | 1380 | The Mammary Gland Undergoes Cycles of Development and Regression | 1426 |
| Interactions Between Epithelium and Mesenchyme Generate Branching Tubular Structures | 1381 | <i>Summary</i> | 1428 |
| <i>Summary</i> | 1382 | SENSORY EPITHELIA | 1429 |
| NEURAL DEVELOPMENT | 1383 | Olfactory Sensory Neurons Are Continually Replaced | 1429 |
| Neurons Are Assigned Different Characters According to the Time and Place Where They Are Born | 1383 | Auditory Hair Cells Have to Last a Lifetime | 1430 |
| The Character Assigned to a Neuron at Its Birth Governs the Connections It Will Form | 1385 | Most Permanent Cells Renew Their Parts: the Photoreceptor Cells of the Retina | 1432 |
| Each Axon or Dendrite Extends by Means of a Growth Cone at Its Tip | 1386 | <i>Summary</i> | 1433 |
| The Growth Cone Pilots the Developing Neurite Along a Precisely Defined Path <i>In Vivo</i> | 1387 | THE AIRWAYS AND THE GUT | 1434 |
| Growth Cones Can Change Their Sensibilities as They Travel | 1389 | Adjacent Cell Types Collaborate in the Alveoli of the Lungs | 1434 |
| Target Tissues Release Neurotrophic Factors That Control Nerve Cell Growth and Survival | 1389 | Goblet Cells, Ciliated Cells, and Macrophages Collaborate to Keep the Airways Clean | 1434 |
| Neuronal Specificity Guides the Formation of Orderly Neural Maps | 1391 | The Lining of the Small Intestine Renews Itself Faster Than Any Other Tissue | 1436 |
| Axons From Different Regions of the Retina Respond Differently to a Gradient of Repulsive Molecules in the Tectum | 1392 | Wnt Signaling Maintains the Gut Stem-Cell Compartment | 1438 |
| Diffuse Patterns of Synaptic Connections Are Sharpened by Activity-Dependent Remodeling | 1393 | Notch Signaling Controls Gut Cell Diversification | 1439 |
| Experience Molds the Pattern of Synaptic Connections in the Brain | 1395 | Ephrin-Eph Signaling Controls the Migrations of Gut Epithelial Cells | 1440 |
| Adult Memory and Developmental Synapse Remodeling May Depend on Similar Mechanisms | 1396 | Wnt, Hedgehog, PDGF, and BMP Signaling Pathways Combine to Delimit the Stem-Cell Niche | 1441 |
| <i>Summary</i> | 1397 | The Liver Functions as an Interface Between the Digestive Tract and the Blood | 1442 |
| PLANT DEVELOPMENT | 1398 | Liver Cell Loss Stimulates Liver Cell Proliferation | 1443 |
| <i>Arabidopsis</i> Serves as a Model Organism for Plant Molecular Genetics | 1398 | Tissue Renewal Does Not Have to Depend on Stem Cells: Insulin-Secreting Cells in the Pancreas | 1444 |
| The <i>Arabidopsis</i> Genome Is Rich in Developmental Control Genes | 1399 | <i>Summary</i> | 1445 |
| Embryonic Development Starts by Establishing a Root-Shoot Axis and Then Halts Inside the Seed | 1400 | BLOOD VESSELS, LYMPHATICS, AND ENDOTHELIAL CELLS | 1445 |
| The Parts of a Plant Are Generated Sequentially by Meristems | 1403 | Endothelial Cells Line All Blood Vessels and Lymphatics | 1445 |
| Development of the Seedling Depends on Environmental Signals | 1403 | Endothelial Tip Cells Pioneer Angiogenesis | 1446 |
| Long-Range Hormonal Signals Coordinate Developmental Events in Separate Parts of the Plant | 1403 | Different Types of Endothelial Cells Form Different Types of Vessel | 1447 |
| The Shaping of Each New Structure Depends on Oriented Cell Division and Expansion | 1406 | Tissues Requiring a Blood Supply Release VEGF; Notch Signaling Between Endothelial Cells Regulates the Response | 1448 |
| Each Plant Module Grows From a Microscopic Set of Primordia in a Meristem | 1407 | Signals from Endothelial Cells Control Recruitment of Pericytes and Smooth Muscle Cells to Form the Vessel Wall | 1450 |
| Polarized Auxin Transport Controls the Pattern of Primordia in the Meristem | 1408 | <i>Summary</i> | 1450 |
| Cell Signaling Maintains the Meristem | 1409 | RENEWAL BY MULTIPOTENT STEM CELLS: BLOOD CELL FORMATION | 1450 |
| Regulatory Mutations Can Transform Plant Topology by Altering Cell Behavior in the Meristem | 1410 | The Three Main Categories of White Blood Cells Are Granulocytes, Monocytes, and Lymphocytes | 1451 |
| The Switch to Flowering Depends on Past and Present Environmental Cues | 1412 | The Production of Each Type of Blood Cell in the Bone Marrow Is Individually Controlled | 1453 |
| Homeotic Selector Genes Specify the Parts of a Flower | 1413 | Bone Marrow Contains Hemopoietic Stem Cells | 1454 |
| <i>Summary</i> | 1415 | A Multipotent Stem Cell Gives Rise to All Classes of Blood Cells | 1456 |
| <i>References</i> | 1415 | Commitment Is a Stepwise Process | 1456 |
| Chapter 23 Specialized Tissues, Stem Cells, and Tissue Renewal | 1417 | Divisions of Committed Progenitor Cells Amplify the Number of Specialized Blood Cells | 1457 |
| EPIDERMIS AND ITS RENEWAL BY STEM CELLS | 1417 | Stem Cells Depend on Contact Signals From Stromal Cells | 1458 |
| Epidermal Cells Form a Multilayered Waterproof Barrier | 1419 | Factors That Regulate Hemopoiesis Can Be Analyzed in Culture | 1459 |
| Differentiating Epidermal Cells Express a Sequence of Different Genes as They Mature | 1420 | Erythropoiesis Depends on the Hormone Erythropoietin | 1459 |
| Stem Cells in the Basal Layer Provide for Renewal of the Epidermis | 1420 | Multiple CSFs Influence Neutrophil and Macrophage Production | 1460 |
| The Two Daughters of a Stem Cell Do Not Always Have to Become Different | 1421 | The Behavior of a Hemopoietic Cell Depends Partly on Chance | 1461 |
| | | Regulation of Cell Survival Is as Important as Regulation of Cell Proliferation | 1462 |
| | | <i>Summary</i> | 1462 |
| | | GENESIS, MODULATION, AND REGENERATION OF SKELETAL MUSCLE | 1463 |
| | | Myoblasts Fuse to Form New Skeletal Muscle Fibers | 1464 |

| | | | |
|--|------|--|------|
| Muscle Cells Can Vary Their Properties by Changing the Protein Isoforms They Contain | 1465 | Pathogens Evolve Rapidly | 1518 |
| Skeletal Muscle Fibers Secrete Myostatin to Limit Their Own Growth | 1465 | Antigenic Variation in Pathogens Occurs by Multiple Mechanisms | 1519 |
| Some Myoblasts Persist as Quiescent Stem Cells in the Adult | 1466 | Error-Prone Replication Dominates Viral Evolution | 1520 |
| <i>Summary</i> | 1467 | Drug-Resistant Pathogens Are a Growing Problem | 1521 |
| FIBROBLASTS AND THEIR TRANSFORMATIONS: THE CONNECTIVE-TISSUE CELL FAMILY | 1467 | <i>Summary</i> | 1524 |
| Fibroblasts Change Their Character in Response to Chemical Signals | 1467 | BARRIERS TO INFECTION AND THE INNATE IMMUNE SYSTEM | 1524 |
| The Extracellular Matrix May Influence Connective-Tissue Cell Differentiation by Affecting Cell Shape and Attachment | 1468 | Epithelial Surfaces and Defensins Help Prevent Infection | 1525 |
| Osteoblasts Make Bone Matrix | 1469 | Human Cells Recognize Conserved Features of Pathogens | 1526 |
| Most Bones Are Built Around Cartilage Models | 1470 | Complement Activation Targets Pathogens for Phagocytosis or Lysis | 1528 |
| Bone Is Continually Remodeled by the Cells Within It | 1472 | Toll-like Proteins and NOD Proteins Are an Ancient Family of Pattern Recognition Receptors | 1530 |
| Osteoclasts Are Controlled by Signals From Osteoblasts | 1473 | Phagocytic Cells Seek, Engulf, and Destroy Pathogens | 1531 |
| Fat Cells Can Develop From Fibroblasts | 1474 | Activated Macrophages Contribute to the Inflammatory Response at Sites of Infection | 1533 |
| Leptin Secreted by Fat Cells Provides Feedback to Regulate Eating | 1475 | Virus-Infected Cells Take Drastic Measures to Prevent Viral Replication | 1534 |
| <i>Summary</i> | 1476 | Natural Killer Cells Induce Virus-Infected Cells to Kill Themselves | 1535 |
| STEM-CELL ENGINEERING | 1476 | Dendritic Cells Provide the Link Between the Innate and Adaptive Immune Systems | 1536 |
| Hemopoietic Stem Cells Can Be Used to Replace Diseased Blood Cells with Healthy Ones | 1477 | <i>Summary</i> | 1537 |
| Epidermal Stem Cell Populations Can Be Expanded in Culture for Tissue Repair | 1477 | <i>References</i> | 1537 |
| Neural Stem Cells Can Be Manipulated in Culture | 1478 | Chapter 25 The Adaptive Immune System | 1539 |
| Neural Stem Cells Can Repopulate the Central Nervous System | 1478 | LYMPHOCYTES AND THE CELLULAR BASIS OF ADAPTIVE IMMUNITY | 1540 |
| Stem Cells in the Adult Body Are Tissue-Specific | 1479 | Lymphocytes Are Required for Adaptive Immunity | 1540 |
| ES Cells Can Make Any Part of the Body | 1480 | The Innate and Adaptive Immune Systems Work Together | 1541 |
| Patient-Specific ES Cells Could Solve the Problem of Immune Rejection | 1481 | B Lymphocytes Develop in the Bone Marrow; T Lymphocytes Develop in the Thymus | 1543 |
| ES Cells Are Useful for Drug Discovery and Analysis of Disease | 1482 | The Adaptive Immune System Works by Clonal Selection | 1544 |
| <i>Summary</i> | 1482 | Most Antigens Activate Many Different Lymphocyte Clones | 1545 |
| <i>References</i> | 1483 | Immunological Memory Involves Both Clonal Expansion and Lymphocyte Differentiation | 1545 |
| Chapter 24 Pathogens, Infection, and Innate Immunity | 1485 | Immunological Tolerance Ensures That Self Antigens Are Not Normally Attacked | 1547 |
| INTRODUCTION TO PATHOGENS | 1486 | Lymphocytes Continuously Circulate Through Peripheral Lymphoid Organs | 1549 |
| Pathogens Have Evolved Specific Mechanisms for Interacting with Their Hosts | 1486 | <i>Summary</i> | 1551 |
| The Signs and Symptoms of Infection May Be Caused by the Pathogen or by the Host's Responses | 1487 | B CELLS AND ANTIBODIES | 1551 |
| Pathogens Are Phylogenetically Diverse | 1488 | B Cells Make Antibodies as Both Cell-Surface Antigen Receptors and Secreted Proteins | 1552 |
| Bacterial Pathogens Carry Specialized Virulence Genes | 1489 | A Typical Antibody Has Two Identical Antigen-Binding Sites | 1552 |
| Fungal and Protozoan Parasites Have Complex Life Cycles with Multiple Forms | 1494 | An Antibody Molecule Is Composed of Heavy and Light Chains | 1552 |
| All Aspects of Viral Propagation Depend on Host Cell Machinery | 1496 | There Are Five Classes of Antibody Heavy Chains, Each with Different Biological Properties | 1553 |
| Prions Are Infectious Proteins | 1498 | The Strength of an Antibody-Antigen Interaction Depends on Both the Number and the Affinity of the Antigen-Binding Sites | 1557 |
| Infectious Disease Agents Are Linked To Cancer, Heart Disease, and Other Chronic Illnesses | 1499 | Antibody Light and Heavy Chains Consist of Constant and Variable Regions | 1558 |
| <i>Summary</i> | 1501 | The Light and Heavy Chains Are Composed of Repeating Ig Domains | 1559 |
| CELL BIOLOGY OF INFECTION | 1501 | An Antigen-Binding Site Is Constructed from Hypervariable Loops | 1560 |
| Pathogens Cross Protective Barriers to Colonize the Host | 1501 | <i>Summary</i> | 1561 |
| Pathogens That Colonize Epithelia Must Avoid Clearance by the Host | 1502 | THE GENERATION OF ANTIBODY DIVERSITY | 1562 |
| Intracellular Pathogens Have Mechanisms for Both Entering and Leaving Host Cells | 1504 | Antibody Genes Are Assembled From Separate Gene Segments During B Cell Development | 1562 |
| Virus Particles Bind to Molecules Displayed on the Host Cell Surface | 1505 | Imprecise Joining of Gene Segments Greatly Increases the Diversity of V Regions | 1564 |
| Virions Enter Host Cells by Membrane Fusion, Pore Formation, or Membrane Disruption | 1506 | The Control of V(D)J Recombination Ensures That B Cells Are Monospecific | 1565 |
| Bacteria Enter Host Cells by Phagocytosis | 1507 | Antigen-Driven Somatic Hypermutation Fine-Tunes Antibody Responses | 1566 |
| Intracellular Eucaryotic Parasites Actively Invade Host Cells | 1508 | B Cells Can Switch the Class of Antibody They Make | 1567 |
| Many Pathogens Alter Membrane Traffic in the Host Cell | 1511 | <i>Summary</i> | 1568 |
| Viruses and Bacteria Use the Host Cell Cytoskeleton for Intracellular Movement | 1514 | | |
| Viral Infections Take Over the Metabolism of the Host Cell | 1517 | | |
| Pathogens Can Alter the Behavior of the Host Organism to Facilitate the Spread of the Pathogen | 1518 | | |

| | | | |
|--|------|---|------|
| T CELLS AND MHC PROTEINS | 1569 | | |
| T Cell Receptors (TCRs) Are Antibodylike Heterodimers | 1570 | Most Developing Cytotoxic and Helper T Cells That Could Be Activated by Self-Peptide–MHC Complexes Are Eliminated in the Thymus | 1586 |
| Antigen Presentation by Dendritic Cells Can Either Activate or Tolerize T Cells | 1571 | Some Organ-Specific Proteins Are Ectopically Expressed in the Thymus Medulla | 1587 |
| Effector Cytotoxic T Cells Induce Infected Target Cells to Kill Themselves | 1572 | The Function of MHC Proteins Helps Explain Their Polymorphism | 1588 |
| Effector Helper T Cells Help Activate Other Cells of the Innate and Adaptive Immune Systems | 1573 | <i>Summary</i> | 1588 |
| Regulatory T Cells Suppress the Activity of Other T Cells | 1574 | HELPER T CELLS AND LYMPHOCYTE ACTIVATION | 1589 |
| T Cells Recognize Foreign Peptides Bound to MHC Proteins | 1575 | Activated Dendritic Cells Use Multiple Mechanisms to Activate T Cells | 1590 |
| MHC Proteins Were Identified in Transplantation Reactions Before Their Functions Were Known | 1575 | The Activation of T Cells Is Controlled by Negative Feedback | 1591 |
| Class I and Class II MHC Proteins Are Structurally Similar Heterodimers | 1576 | The Subclass of Effector Helper T Cell Determines the Nature of the Adaptive Immune Response | 1592 |
| An MHC Protein Binds a Peptide and Interacts with a T Cell Receptor | 1577 | T _H 1 Cells Activate Infected Macrophages and Stimulate An Inflammatory Response | 1594 |
| MHC Proteins Help Direct T Cells to Their Appropriate Targets | 1579 | Antigen Binding to B Cell Receptors (BCRs) Is Only One Step in B Cell Activation | 1595 |
| CD4 and CD8 Co-Receptors Bind to Invariant Parts of MHC Proteins | 1580 | Antigen-Specific Helper T Cells Are Essential for Activating Most B Cells | 1597 |
| Cytotoxic T Cells Recognize Fragments of Foreign Cytosolic Proteins in Association with Class I MHC Proteins | 1581 | A Special Class of B Cells Recognize T-Cell-Independent Antigens | 1598 |
| Helper T Cells Respond to Fragments of Endocytosed Foreign Protein Associated with Class II MHC Proteins | 1583 | Immune Recognition Molecules Belong to the Ancient Ig Superfamily | 1599 |
| Potentially Useful T Cells Are Positively Selected in the Thymus | 1585 | <i>Summary</i> | 1600 |
| | | <i>References</i> | 1600 |