Sonderheft 297 Special Issue



Landbauforschung Völkenrode FAL Agricultural Research

Investigations on possibilities to improve the antiphytopathogenic potential of soils against the cyst nematode *Heterodera schachtii* and the citrus nematode *Tylenchulus semipenetrans*

Hazem Abdelnabby

ULB Darmstadt

TABLE OF CONTENTS

Table of contents	I
List of abbreviations	III
List of tables	IV
List of figures	VI
1 Introduction	1
1.1 Background.	1
1.2 Objectives	5
2 Materials and methods	6
2.1 Sampling procedures	6
2.2 Biological methods	6
2.2.1 Nematode propagation in soil	6
2.2.2 Inoculation procedures	7
2.2.3 Nematode extraction, counting and identification	7
2.3 Vegetative Measurements	8
2.4 Nematode control strategies	8
2.4.1 Organic and inorganic fertilisers	8
2.4.2 Organic and amino acids	8
2.4.3 Microbial agents	9
2.4.4 Navel orange on resistant rootstocks	9
2.4.5 Plant extracts	10
2.4.6 Carbofuran	12
2.5 Experimental design	12
2.5.1 Greenhouse experiment of cyst nematode control strategy - CYNO	COS12
Enhancing the antiphytopathogenic potential of the soil against cyst ner	natodes 12
2.5.2 Greenhouse experiment of the citrus nematode control strategy - C	CINCOS15
Enhancing the antiphytopathogenic potential of the soil against citrus n	ematodes 15
2.5.3 Greenhouse experiment of citrus rootstocks susceptibility to the in	fection
with citrus nematodes - ROSSUS	16
Application of citrus rootstocks as citrus nematode control strategy	16
2.5.4 Field experiment with citrus trees – FECIT	17
Timing of application	17
2.6 Statistical analysis	18

3 Results	19
3.1 Enhancing the antiphytopathogenic potential of soil against the sugar beet cyst	
nematode Heterodera schachtii	19
3.1.1 Organic and inorganic fertilisers	19
3.1.2 Microbial agents	24
3.1.3 Plant extracts	28
3.2 Enhancing the antiphytopathogenic potential of soil against the citrus nematode	
Tylenchulus semipenetrans	38
3.2.1 Organic and inorganic fertilisers	38
3.2.2 Organic and amino acids	42
3.2.3 Microbial agents	45
3.2.4 Plant extracts	50
3.3 Application of different rootstocks as a mean to control citrus nematode	58
3.3.1 Relative susceptibility of citrus rootstocks	58
3.3.2 Application of Navel orange scions on resistant rootstocks against citrus	
nematode	63
3.4 The suitable application time of nematode control strategies	67
4 Discussion	69
4.1 Evaluation of the approaches used to enhance the antiphytopathogenic potential	
of soil against the sugar beet cyst nematode Heterodera schachtii and the citrus	
nematode Tylenchulus semipenetrans	71
4.2 Evaluation of different rootstocks as a mean to control citrus nematodes	83
4.3 Determination of the most suitable application time for nematode control	
strategies	84
5 Summary/ Zusammenfassung	86
6 References	92
7 Appendix	05